

Stochastic Models: Exercise 5

1. Let $\{X_n : n \geq 0\}$ be an irreducible Markov chain with period $d \geq 1$ thus for any state i

$$d = \gcd\{n \geq 1 : P[X_n = i \mid X_0 = i] > 0\}$$

Suppose $\{X_{nd} : n \geq 0\}$ is not aperiodic, thus for some integer $k > 1$, we have

$$k = \gcd\{n \geq 1 : P[X_{nd} = i \mid X_0 = i] > 0\}$$

for every state i . This implies that, for all states, the number of transitions needed to return to state i given that it starts from i in $\{X_n : n \geq 0\}$ is of the form lkd where l is a positive integer. This contradicts that $\{X_n : n \geq 0\}$ is a Markov chain with period d since for any integer l , $lkd > d$, thus $k = 1$ as required and $\{X_{nd} : n \geq 0\}$ as aperiodic.

If we consider the states accessible to $\{X_{nd} : n \geq 0\}$, it is irreducible. Else it is not. Consider the simple symmetric random walk where $\{X_{2n} : n \geq 0\}$ is aperiodic and irreducible for the even states. But if we consider both even and odd states, it is not irreducible as it cannot visit an odd state in even number of steps.

2. Suppose $i \leftrightarrow j$ and let i be positive recurrent, thus $\sum_{n=1}^{\infty} n f_{ii}^n < \infty$.

$$\infty > \sum_{n=1}^{\infty} n f_{ii}^n \geq \sum_{n=1}^{\infty} n f_{ji} f_{ii}^n f_{ij} \geq \sum_{n=1}^{\infty} n f_{jj}^n$$

3. Let $\{X_n : n \geq 0\}$ be an irreducible and aperiodic Markov chain. The chain is doubly stochastic, thus $\sum_i P_{ij} = 1$. For any two states i and j , we have $i \leftrightarrow j$ since the Markov chain is irreducible and together with the aperiodicity, we have $\lim_{n \rightarrow \infty} P_{ij}^n = \pi_j$.

$$\begin{aligned} \pi_j &= \lim_{n \rightarrow \infty} P_{ij}^{n+1} = \lim_{n \rightarrow \infty} \sum_{l=0}^k P_{il}^n P_{lj} \\ &= \sum_{l=0}^k \lim_{n \rightarrow \infty} P_{il}^n P_{lj} \\ &= \sum_{l=0}^k \pi_l P_{lj} \end{aligned}$$

by assuming the uniform distribution, show that the above is satisfied.

4. Let $\{X_n : n \geq 0\}$ be a Markov chain with states $S = \{0, 1, 2, 3, 4\}$ denoting the number of umbrella(s) in his home. The transition matrix \mathbb{P} is such that $P_{i,i-1} = p = 1 - P_{i,i}$ for $i \neq 0$ and $P_{0,0} = 1$.

(a)

(b)

5. (a)

(b)

6. (a)

(b)