

Statistics: Homework 2

6.3 Given $\hat{\theta} = 2\bar{X}_n$ and $X_1, \dots, X_n \sim \text{Uniform}(0, \theta)$,

$$\begin{aligned}\text{bias}(\hat{\theta}) &= \mathbb{E}(2\bar{X}_n) - \theta \\ &= 2n^{-1} \mathbb{E} \left(\sum_{i=1}^n X_i \right) - \theta \\ &= 2n^{-1} \sum_{i=1}^n \mathbb{E}(X_i) - \theta \\ &= 2n^{-1} \frac{n\theta}{2} - \theta = 0 \\ \text{se}(\hat{\theta})^2 &= \mathbb{V}(2\bar{X}_n) \\ &= 4\mathbb{V}(\bar{X}_n) \\ &= 4n^{-2} \mathbb{V} \left(\sum_{i=1}^n X_i \right) \\ &= 4n^{-2} \sum_{i=1}^n \mathbb{V}(X_i) \\ &= 4n^{-2} \frac{n\theta^2}{12} = \frac{\theta^2}{3n} \\ \text{MSE}(\hat{\theta}) &= \text{bias}(\hat{\theta})^2 + \text{se}(\hat{\theta})^2 = \frac{\theta^2}{3n}\end{aligned}$$

7.2 For $X_1, \dots, X_n \sim \text{Bernoulli}(p)$ plug-in estimator for p is

$$\hat{p} = \frac{1}{n} \sum_{i=1}^n X_i$$

and the estimated standard error is given by

$$\hat{\text{se}}_p = \sqrt{\mathbb{V}(\hat{p})} = \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}$$

As the X_i 's are iid, by Central Limit Theorem, \hat{p} is asymptotically normal with mean p and variance $\hat{\text{se}}_p^2$. Thus an approximate 90% confidence interval for p is $(\hat{p} - 1.96\hat{\text{se}}_p, \hat{p} + 1.96\hat{\text{se}}_p)$.

For $X_1, \dots, X_n \sim \text{Bernoulli}(p)$ and $Y_1, \dots, Y_m \sim \text{Bernoulli}(q)$ plug-in estimator for $p - q$ is

$$\hat{p} - \hat{q} = \frac{1}{n} \sum_{i=1}^n X_i - \frac{1}{m} \sum_{i=1}^m Y_i$$

with estimated standard error

$$\hat{\text{se}}_{p-q} = \sqrt{\mathbb{V}(\hat{p} - \hat{q})} = \sqrt{\mathbb{V}(\hat{p}) + \mathbb{V}(\hat{q})} = \sqrt{\frac{\hat{p}(1 - \hat{p})}{n} + \frac{\hat{q}(1 - \hat{q})}{m}}$$

Since the Y_i 's are iid, by Central Limit Theorem \hat{q} is asymptotically normal with mean q and variance $\hat{\text{se}}_q^2$. The difference of two asymptotically normal random variables is asymptotically normal, thus $\hat{p} - \hat{q}$ is asymptotically normal with mean $p - q$ and variance $\hat{\text{se}}_{p-q}^2$. An approximate 90% confidence interval is

$$\text{se}_{p-q} = (\hat{p} - \hat{q} - 1.96\hat{\text{se}}_{p-q}, \hat{p} - \hat{q} + 1.96\hat{\text{se}}_{p-q})$$

7.9 An estimate for $p_1 - p_2$ is $0.9 - 0.85 = 0.05$ with standard error

$$\sqrt{\frac{0.9(1 - 0.9)}{100} + \frac{0.85(1 - 0.85)}{100}} = 0.0466368953$$

with 80% and 90% confidence intervals given by

$$80\% : (\hat{p} - \hat{q} - 1.282\text{se}_{p-q}, \hat{p} - \hat{q} + 1.282\text{se}_{p-q}) = (-0.0097885, 0.1097885)$$

$$90\% : (\hat{p} - \hat{q} - 1.96\text{se}_{p-q}, \hat{p} - \hat{q} + 1.96\text{se}_{p-q}) = (-0.041408315, 0.141408315)$$

8.7

9.2

9.6