

## Statistics: Homework 2

6.3 Given  $\hat{\theta} = 2\bar{X}_n$  and  $X_1, \dots, X_n \sim \text{Uniform}(0, \theta)$ ,

$$\begin{aligned}\text{bias}(\hat{\theta}) &= \mathbb{E}(2\bar{X}_n) - \theta \\ &= 2n^{-1} \mathbb{E} \left( \sum_{i=1}^n X_i \right) - \theta \\ &= 2n^{-1} \sum_{i=1}^n \mathbb{E}(X_i) - \theta \\ &= 2n^{-1} \frac{n\theta}{2} - \theta = 0 \\ \text{se}(\hat{\theta})^2 &= \mathbb{V}(2\bar{X}_n) \\ &= 4\mathbb{V}(\bar{X}_n) \\ &= 4n^{-2} \mathbb{V} \left( \sum_{i=1}^n X_i \right) \\ &= 4n^{-2} \sum_{i=1}^n \mathbb{V}(X_i) \\ &= 4n^{-2} \frac{n\theta^2}{12} = \frac{\theta^2}{3n} \\ \text{MSE}(\hat{\theta}) &= \text{bias}(\hat{\theta})^2 + \text{se}(\hat{\theta})^2 = \frac{\theta^2}{3n}\end{aligned}$$

7.2 For  $X_1, \dots, X_n \sim \text{Bernoulli}(p)$  plug-in estimator for  $p$  is

$$\hat{p} = \frac{1}{n} \sum_{i=1}^n X_i$$

and the estimated standard error is given by

$$\hat{\text{se}}_p = \sqrt{\mathbb{V}(\hat{p})} = \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}$$

As the  $X_i$ 's are iid, by Central Limit Theorem,  $\hat{p}$  is asymptotically normal with mean  $p$  and variance  $\hat{\text{se}}_p^2$ . Thus an approximate 90% confidence interval for  $p$  is  $(\hat{p} - 1.645\hat{\text{se}}_p, \hat{p} + 1.645\hat{\text{se}}_p)$ .

For  $X_1, \dots, X_n \sim \text{Bernoulli}(p)$  and  $Y_1, \dots, Y_m \sim \text{Bernoulli}(q)$  plug-in estimator for  $p - q$  is

$$\hat{p} - \hat{q} = \frac{1}{n} \sum_{i=1}^n X_i - \frac{1}{m} \sum_{i=1}^m Y_i$$

with estimated standard error

$$\hat{\text{se}}_{p-q} = \sqrt{\mathbb{V}(\hat{p} - \hat{q})} = \sqrt{\mathbb{V}(\hat{p}) + \mathbb{V}(\hat{q})} = \sqrt{\frac{\hat{p}(1 - \hat{p})}{n} + \frac{\hat{q}(1 - \hat{q})}{m}}$$

Since the  $Y_i$ 's are iid, by Central Limit Theorem  $\hat{q}$  is asymptotically normal with mean  $q$  and variance  $\hat{\text{se}}_q^2$ . The difference of two asymptotically normal random variables is asymptotically normal, thus  $\hat{p} - \hat{q}$  is asymptotically normal with mean  $p - q$  and variance  $\hat{\text{se}}_{p-q}^2$ . An approximate 90% confidence interval is

$$(\hat{p} - \hat{q} - 1.645\hat{\text{se}}_{p-q}, \hat{p} - \hat{q} + 1.645\hat{\text{se}}_{p-q})$$

7.9 An estimate for  $p_1 - p_2$  is  $0.9 - 0.85 = 0.05$  with standard error

$$\sqrt{\frac{0.9(1 - 0.9)}{100} + \frac{0.85(1 - 0.85)}{100}} = 0.0466368953$$

with 80% and 90% confidence intervals given by

$$\begin{aligned} 80\% : \quad & (\hat{p} - \hat{q} - 1.282\text{se}_{p-q}, \hat{p} - \hat{q} + 1.282\text{se}_{p-q}) = (-0.0097885, 0.1097885) \\ 90\% : \quad & (\hat{p} - \hat{q} - 1.96\text{se}_{p-q}, \hat{p} - \hat{q} + 1.96\text{se}_{p-q}) = (-0.041408315, 0.141408315) \end{aligned}$$

8.7 (a)

$$\begin{aligned} \mathbb{P}(\hat{\theta} \leq k) &= \mathbb{P}(\max\{X_1, \dots, X_n\} \leq k) \\ &= \prod_{i=1}^n \mathbb{P}(X_i \leq k) \\ &= \left(\frac{k}{\theta}\right)^n \end{aligned}$$




Figure 1: Comparison of the true distribution  $\hat{\theta}$  to histograms from bootstrap

(b) Let  $\hat{\theta} = X_{max} = \max\{X_1, \dots, X_n\}$ . Then

$$\begin{aligned} \mathbb{P}(\hat{\theta}^* = \hat{\theta}) &= 1 - \mathbb{P}(\hat{\theta}^* \neq \hat{\theta}) \\ &= 1 - \left(1 - \frac{1}{n}\right)^n \end{aligned}$$

The second equality holds as  $\mathbb{P}(\hat{\theta}^* \neq \hat{\theta})$  denotes the probability that any random sampling with replacement of the  $n$  samples drawn has probability of  $1 - 1/n$  not being  $x_{max}$  (which is fixed since a random sample of  $n$  has been drawn). As each sampling process is iid due to replacement, probability of them all not being  $x_{max}$  is  $(1 - 1/n)^n$ . Thus we have  $\mathbb{P}(\hat{\theta}^* = \hat{\theta}) \rightarrow 0$  as  $n \rightarrow \infty$  and  $\mathbb{P}(\hat{\theta}^* = \hat{\theta}) = .632$  for  $n = 50$ .

9.2 (a) For  $X \sim \text{Uniform}(a, b)$

$$\begin{aligned} \hat{\mu}\bar{X}_n &= \frac{1}{n} \sum_{i=1}^n X_i = \frac{\hat{b} + \hat{a}}{2} \\ \hat{\sigma}^2 \frac{1}{n} \sum_{i=1}^n (\bar{X}_n - X_i)^2 &= \frac{(b-a)^2}{12} \end{aligned}$$

thus

$$\begin{aligned}(b-a) + (b+a) &= +\sqrt{12\hat{\sigma}^2} + 2\hat{\mu} \\ b &= \frac{1}{2} \left( \sqrt{12\hat{\sigma}^2} + 2\hat{\mu} \right) \\ a &= 2\hat{\mu} - b\end{aligned}$$

the positive root is taken as  $b - a > 0$ .

(b) The log-likelihood function is given by

$$\ell_n(a, b) = \sum_{i=1}^n \log \frac{X_i}{b-a}$$

(c)

(d)

9.6