

Real Analysis: Homework 1

1. (a) \mathbb{R} is second-countable by considering the countable basis

$$\mathcal{B} := \{(a, b) | a < b, a, b \in \mathbb{Q}\}$$

We now claim that $\mathcal{B}^n = \{U_1 \times \dots \times U_n | \text{each } U_i \in \mathcal{B} \text{ for } i = 1, \dots, n\}$ is a countable basis for \mathbb{R}^n . It is clear that \mathcal{B}^n is countable as the Cartesian product of countable sets is still countable. To show \mathcal{B}^n is a basis for \mathbb{R}^n :

- (1) Pick $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ and consider the projection map $\pi_i : \mathbb{R}^n \rightarrow \mathbb{R}, (x_1, \dots, x_n) \mapsto x_i$. Thus for each $\pi_i(x) = x_i$ we can find $B_i \in \mathcal{B}$ such that $x_i \in B_i$. Thus $B_1 \times \dots \times B_n$ is the basis element in \mathcal{B}^n containing x .
- (2) Let x belong to the intersection of two basis elements $U = B_1 \times \dots \times B_n, U' = B'_1 \times \dots \times B'_n$. Using the projection map, $\pi_i(U) = B_i, \pi_i(U') = B'_i$ and thus there is a basis element $A_i \subseteq B_i \cap B'_i$ for some $A_i \in \mathcal{B}$. Thus $A = A_1 \times \dots \times A_n$ is the basis element in \mathcal{B}^n such that $A \subseteq U \cap U'$.

Thus we have shown that \mathcal{B}^n is a countable basis for \mathbb{R}^n .

- (b) Let U be an open set of \mathbb{R} . If U is a union of countably many open sets we can simply pick the disjoint open intervals from that union and we are done. Suppose U is an uncountable union of open sets, and without loss of generality assume that they are disjoint, $U = \sqcup_{\alpha \in A} V_\alpha$ for uncountable A , then since \mathbb{R} is second-countable, there exists a countable basis \mathcal{B} for \mathbb{R} . Thus for each $x \in V_\alpha$, there exists $B_x \in \mathcal{B}$ such that $x \in B_x \subseteq V_\alpha$. Consider the set

$$\left\{ \bigcup_{x \in V_\alpha} B_x \text{ for all } \alpha \in A \right\}$$

we claim that this set is at most countable and is a disjoint set of open intervals. Every element of the set is open since it is a union of $B \in \mathcal{B}$. It is disjoint since by construction $V_\alpha = \bigcup_{x \in V_\alpha} B_x$. Lastly, suppose that the set above is uncountable, then since there are countably many $B \in \mathcal{B}$, $\left(\bigcup_{x \in V_{\alpha_1}} B_x \right) \cap \left(\bigcup_{x \in V_{\alpha_2}} B_x \right) = V_{\alpha_1} \cap V_{\alpha_2} \neq \emptyset$ for some distinct α_1, α_2 which contradicts the earlier assumption.

2. Let $f : (X, \tau_X) \rightarrow (Y, \tau_Y)$ be a continuous function. Let (X, τ'_X) be a finer topology than (X, τ_X) then $\tau'_X \supseteq \tau_X$. Thus for any $U \in \tau_Y$, $f^{-1}(U) \in \tau_X \subseteq \tau'_X$. Thus $f^{-1}(U) \in \tau'_X$ and $f : (X, \tau'_X) \rightarrow (Y, \tau_Y)$ remains continuous. Let (Y, τ'_Y) is a topology coarser than (Y, τ_Y) and so $\tau_Y \supseteq \tau'_Y$. Hence for $U \in \tau'_Y \subseteq \tau_Y$, we have $f^{-1}(U) \in \tau_X$. Thus $f : (X, \tau_X) \rightarrow (Y, \tau'_Y)$ remains continuous.

3. We shall show that $\int_{\mathbb{R}} e^{-x^2/2} dx = \sqrt{2\pi}$. Start by considering its square,

$$\begin{aligned}
\left(\int_{\mathbb{R}} e^{-x^2/2} dx\right)^2 &= \left(\int_{\mathbb{R}} e^{-x^2/2} dx\right) \left(\int_{\mathbb{R}} e^{-y^2/2} dy\right) \\
&= \int_{\mathbb{R}^2} e^{-\frac{x^2+y^2}{2}} dy dx \\
&= \int_0^{2\pi} \int_0^\infty e^{-r^2/2} r dr d\theta \quad \text{by polar change of coordinates.} \\
&= 2\pi \int_0^\infty e^{-r^2/2} r dr \\
&= 2\pi \int_0^\infty \frac{1}{2} e^{-s/2} ds \quad \text{change of coordinates, } s = r^2 \\
&= \pi \left[-2e^{-s/2} \right]_0^\infty = 2\pi
\end{aligned}$$

which shows what is required and hence $\frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-x^2/2} dx = 1$.

4. Let $x_n > 0$ for all n and $x_n \rightarrow a$ with $a > 0$. Since $f(x) = \log x$ is a continuous function, it suffices to show

$$\frac{1}{n^2} \sum_{k=1}^n k \log x_k \rightarrow \frac{1}{2} \log a$$

Consider the sequence $s_n := \frac{1}{n^2} \sum_{i=1}^n i \log x_i$. Then for some positive integer k , there exists N_1 such that

$$s_n = \frac{1}{n^2} \sum_{i=1}^k i \log x_i + \frac{1}{n^2} \sum_{i=k+1}^n i \log x_i \rightarrow \frac{1}{n^2} \sum_{i=k+1}^n i \log x_i$$

for $n \geq N_1$. Since $\log x_n \rightarrow \log a$, $\limsup_{n \rightarrow \infty} \log x_n = \lim_{n \rightarrow \infty} \log x_n = \log a$. Thus there exists N_2 such that we have $\log x_n < \log a + \epsilon$ for $n \geq N_2$ and arbitrary $\epsilon > 0$. Taking $N = \max\{N_1, N_2\}$, for $n \geq N$,

$$s_n \leq \frac{1}{n^2} \sum_{i=k+1}^n i(\log a + \epsilon) = (\log a + \epsilon) \frac{n^2 - k^2 + n + k}{2n^2}$$

Then $\limsup_{n \rightarrow \infty} s_n \leq \limsup_{n \rightarrow \infty} \frac{1}{n^2} \sum_{i=k+1}^n i(\log a + \epsilon) = \frac{1}{2}(\log a + \epsilon)$. In a similar fashion, we have $\liminf_{n \rightarrow \infty} \log x_n = \log a$ and there exists N_3 such that $\log x_n > \log a - \epsilon$ for $n \geq N_3$ and arbitrary $\epsilon > 0$. Taking $M = \max\{N_1, N_3\}$, for $n \geq M$,

$$s_n \geq \frac{1}{n^2} \sum_{i=k+1}^n i(\log a - \epsilon) = (\log a - \epsilon) \frac{n^2 - k^2 + n + k}{2n^2}$$

Then $\liminf_{n \rightarrow \infty} s_n \geq \liminf_{n \rightarrow \infty} \frac{1}{n^2} \sum_{i=k+1}^n i(\log a - \epsilon) = \frac{1}{2}(\log a - \epsilon)$. As ϵ is arbitrarily chosen we have $\limsup_{n \rightarrow \infty} s_n = \frac{1}{2} \log a = \liminf_{n \rightarrow \infty} s_n$, this $s_n \rightarrow \frac{1}{2} \log a$.

5. We first observe that

$$\int_0^1 f(x) dx = \sum_{j=0}^{n-1} \int_{j/n}^{j+1/n} f(x) dx$$

thus

$$\begin{aligned}
\left| \sum_{j=0}^{n-1} \frac{f(j/n)}{n} - \int_0^1 f(x) dx \right| &= \left| \sum_{j=0}^{n-1} \int_{j/n}^{(j+1)/n} f(j/n) - f(x) dx \right| \\
&= \left| \sum_{j=0}^{n-1} \int_{j/n}^{(j+1)/n} (j/n - x) \cdot \frac{f(j/n) - f(x)}{j/n - x} dx \right| \\
&\leq M \left| \sum_{j=0}^{n-1} \int_{j/n}^{(j+1)/n} (j/n - x) dx \right| \text{ by MVT for integrals and then MVT} \\
&= M \left| \sum_{j=0}^{n-1} -\frac{1}{2} \left(\frac{j}{n} - \frac{j+1}{n} \right)^2 \right| \\
&= \frac{M}{2n}
\end{aligned}$$