
Toxicity stress-test cohorts: every headline AUC carries a design caveat
a
GSE189125 · CD4-Tem fraction (PBMC)
n = 13 (6 severe / 7 no)
0.905
no CI in frozen artifacts;
authors' Suppl. Table 3 value;
significant in only 2/5 denominator closures
n = 13 (6 / 7)
GSE189125 · CD4-T proportion S0
0.690
wide CI spans nearly the
full AUC range at n = 13
GSE216329 · S0 12-dim (0.632-corrected)
n = 24 (15 irAE / 9)
0.804
ANTICONSERVATIVE: 0.632-corrected 0.804
EXCEEDS apparent 0.667; endpoint = any irAE
grade 2-3 (organ-specific), NOT grade ≥3 severe
GSE253720 · CPI-colitis vs HC (probe)
n = 12 (9 CPI / 3 HC)
0.919
case-control confound (HC comparator
without ICI exposure); biopsies at ACTIVE
steroid-experienced/refractory colitis —
not baseline; upper CI unclipped (>1)
Pooled DL (k = 2, descriptive)
k = 2 cohorts
0.777
inherits the anticonservative
0.804 input; descriptive only
0.2
0.4
0.6
0.8
1.0
1.2
AUC (toxicity)
diamond = point estimate without a frozen CI; pooled row = DerSimonian-Laird
over the two pooled-eligible cohorts (descriptive only, k = 2).
Sources: data/v60_tox_meta.json; data/v74/v74_joint_stress_test.json
(toxicity_cohort_recheck); data/v82/v82_provenance_ledger.json (endpoint definitions).