

Chapter 25.3 Part I: Homologous Recombination

1. Homologous recombination in bacteria
 - Major role at stalled replication forks
 - Machinery: central role of RecA
 - Replication restart
2. Homologous recombination in eukaryotes
 - Role in meiosis

Homologous Recombination

Genetic exchanges between molecules that
share sequence similarity

- In principle, can occur at any sequence
- Bacteria: primarily repair function, can also occur during conjugation
- Eukaryotes: repair and meiosis

Homologous recombination in bacteria

Primary function: repair of stalled replication forks




Figure 25-29
Lehninger Principles of Biochemistry, Seventh Edition
© 2017 W. H. Freeman and Company

Homologous recombination in bacteria






Figure 25-30
Lehninger Principles of Biochemistry, Seventh Edition

Homologous recombination in bacteria

Step 1:



RecB and RecD: ATP-dependent helicase

RecB: nuclease activity

RecC: binds to **chi**

RecBCD creates 3'-overhang




Figure 25-31b part 1
Lehninger Principles of Biochemistry, Seventh Edition
© 2017 W. H. Freeman and Company

Chi- 1009 sequences spread around *E. coli* genome





Figure 25-31b part 2
Lehninger Principles of Biochemistry, Seventh Edition
© 2017 W. H. Freeman and Company

Homologous recombination in bacteria

Step 2:

RecBCD



RecA






Figure 25-32c
Lehninger Principles of Biochemistry, Seventh Edition
© 2017 W. H. Freeman and Company

RecA:

- filament on ssDNA
- when interacts with homologous duplex, starts to catalyze strand exchange
- sometimes called "bacterial recombinase"

RecA catalyzes strand exchange



Triple helical structure:
ssDNA (dark strand)
running along the minor
groove of the double helix
to be invaded

Homologous recombination in bacteria

Steps 3 and 4:






Figure 25-30
Lehninger Principles of Biochemistry, Seventh Edition
© 2017 W. H. Freeman and Company

Holliday intermediates can be resolved in two different ways



Stalled replication fork repair- finish the job

Once the fork is reformed- need to restart DNA synthesis



Origin-independent- **replication restart primosome**

- Requires 7 proteins

 - DnaB and helicase loading complex (DnaC)**

 - Primase (DnaG)**

 - specialized helicases (PriABC, DnaT)**

Also requires: Topoisomerases, other nucleases,
polymerases, DNA ligase

Chapter 25.3 Part I: Homologous Recombination

1. Homologous recombination in bacteria
 - Major role at stalled replication forks
 - Machinery: central role of RecA
 - Replication restart
2. Homologous recombination in eukaryotes
 - Role in meiosis

Double strand breaks in eukaryotes

1) Non homologous end joining (NHEJ)

- major pathway in **somatic cells**

requires protein heterodimer (Ku)

Ku70 and Ku80

2) Homologous recombination repair

- when homologous chromosome is nearby: a
programmed role in meiosis

NHEJ in somatic cells




Figure 25-37 part 1
Lehninger Principles of Biochemistry, Seventh Edition
© 2017 W. H. Freeman and Company




Figure 25-37 part 2
Lehninger Principles of Biochemistry, Seventh Edition
© 2017 W. H. Freeman and Company

Homologous recombination during meiosis

"crossing over" reactions at **chiasma**




Figure 25-32
Lehninger Principles of Biochemistry, Fifth Edition
© 2008 W.H. Freeman and Company

although there are "hot spots"- can happen with relatively equal probability along chromosomes- thus could use for gene mapping experiments