

Ch 27.2- Protein synthesis

1) Assembling the machinery and overview

- **Ribosomes:** RNP, structure and properties
- **Aminoacyl-tRNA:** structure and recognition
- Basic mechanisms and architecture

2) A stepwise process

- **Initiation:** steps up to formation of 1st peptide bond
- **Elongation:** synthesis of the 1st bond to addition of the last amino acid
- **Termination:** release completed polypeptide chain
- **Ribosome recycling:** disassembly of the ribosome for next use

Components of bacterial vs eukaryotic ribosome




Figure 27-16
 Lehninger Principles of Biochemistry, Sixth Edition
 © 2013 W. H. Freeman and Company

16S rRNA conserved secondary structures





Figure 27-15
Lehninger Principles of Biochemistry, Sixth Edition
© 2013 W. H. Freeman and Company

Structure of the *T. thermophilus* 70S ribosome



Yusupov (Noller) Science 2001

RNA-only active site





Figure 27-13b
Lehninger Principles of Biochemistry, Fifth Edition
 © 2008 W. H. Freeman and Company

Structure of the *T. thermophilus* 70S ribosome



Yusupov (Noller) Science 2001

General cloverleaf secondary structure of tRNAs




Figure 27-17
Lehninger Principles of Biochemistry, Sixth Edition
© 2013 W. H. Freeman and Company

Cytosolic/bacterial tRNA: archaea and organelles can exhibit differences

Extensive tertiary interactions in tRNA structure




Figure 27-18
Lehninger Principles of Biochemistry, Sixth Edition
© 2013 W. H. Freeman and Company

Aminoacyl-tRNA synthetases (aaRS) charge tRNAs

Nomenclature:

- tRNAs: Arg-tRNA^{Arg}, Lys-tRNA^{Val}
- synthetases: ArgRS, GlyRS, etc

Properties:


- Form an ester-linked amino acid at 3'-end of tRNA
- One aaRS per amino acid (mostly)
- Two classes (Class I and Class II)
 - structurally distinct but same catalytic activity
 - arose independently in evolution

Mechanism:

- Aminoacyl-adenylate
- nucleophilic attack on activated carbonyl

Prototypical aaRS from two different classes

E. coli GlnRS (class I)



Yeast AspRS (Class II)





Figure 27-22
Lehninger Principles of Biochemistry, Sixth Edition
© 2013 W. H. Freeman and Company

The central dogma: fidelity of gene expression



Lewin's Essential Genes

Fidelity of aminoacylation by aaRS

The problem: many similar-looking substrates for aaRS

Use of "identity elements": defined sequences that specify a certain amino acid

- determinants and antideterminants
- test: necessary and sufficient for recognition

"A 2nd genetic code"...

tRNA^{Ala} identity elements- conserved G-U base pair




Figure 27-23
Lehninger Principles of Biochemistry, Sixth Edition
© 2013 W. H. Freeman and Company

Fidelity of aminoacylation by aaRS

The problem: many similar-looking substrates for aaRS

Additional strategies used by some aaRS:

1) Kinetic mechanism:


- aminoacylation is faster with cognate tRNA than non-cognate

2) Chemical proofreading:

- hydrolyze incorrect aa-tRNA combinations


Fidelity of aminoacylation by aaRS

Chemical proofreading (editing)



Valine

Unnumbered 27 p1081
Lehninger Principles of Biochemistry, Fifth Edition
© 2008 W. H. Freeman and Company



Isoleucine

- tRNA not always required
- aaRS-specific
- "Double sieve" mechanism
Ex. IleRS:
 - 1st: size for aa-AMP formation, Leu won't fit
 - 2nd: size for editing site, Val fits, Ile doesn't

Two types of editing (depending on aaRS):

- 1) hydrolyze non-cognate aa-AMP (pre-transfer)
- 2) hydrolyze mischarged aa-tRNA (post-transfer)