

Due: September 9, 2016

MATH 320: HOMEWORK 1

Please read through chapters 2 and 3 of the textbook. Obtain access to a copy of MATLAB or GNU Octave to answer the following questions. For all questions, please write a brief description of how your code works.

- (1) Problem 2.9: The density of freshwater can be computed as a function of temperature with the following cubic equation:

$$\rho = 5.5289 \times 10^{-8} T_C^3 - 8.5016 \times 10^{-6} T_C^2 + 6.5622 \times 10^{-5} T_C + 0.99987$$

where ρ = density (g/cm³) and T_C = temperature (°C). Use MATLAB to generate a vector of temperatures ranging from 32°F to 93.2°F using increments of 3.6°F. Convert this vector to degrees Celsius and then compute a vector of densities based on the cubic formula. Create a plot of ρ versus T_C . Recall that $T_C = 5/9(T_F - 32)$.

- (2) Problem 2.15: The Maclaurin series expansion for the cosine is

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \dots$$

Use MATLAB to create a plot of the cosine (solid line) along with a plot of the series expansion (black dashed line) up to and including the term $x^8/8!$. Use the built-in function `factorial` in computing the series expansion. Make the range of the abscissa from $x = 0$ to $3\pi/2$.

- (3) Problem 3.6: Two distances are required to specify the location of a point relative to an origin in two-dimensional space:

- The horizontal and vertical distances (x, y) in Cartesian coordinates; or
- The radius and angle (r, θ) in polar coordinates.

It is relatively straightforward to compute Cartesian coordinates (x, y) on the basis of polar coordinates (r, θ) . The reverse process is not so simple. The radius can be computed by the following formula: $r = \sqrt{x^2 + y^2}$.

If the coordinates lie within the first and fourth quadrants (i.e., $x > 0$), then a simple formula can be used to compute θ : $\theta = \tan^{-1}(y/x)$. The difficulty arises for the other cases. The following table summarizes the possibilities:

x	y	θ
< 0	> 0	$\tan^{-1}(y/x) + \pi$
< 0	< 0	$\tan^{-1}(y/x) - \pi$
< 0	$= 0$	π
$= 0$	> 0	$\pi/2$
$= 0$	< 0	$-\pi/2$
$= 0$	$= 0$	0

Write a well-structured M-file using `if...elseif` structures to calculate r and θ as a function of x and y . Express the final results for θ in degrees. Test your program by evaluating the following cases:

x	y	r	θ
2	0		
2	1		
0	3		
-3	1		
-2	0		
-1	-2		
0	0		
0	-2		
2	2		

- (4) Problem 3.20: A Cartesian vector can be thought of as representing magnitudes along the x -, y -, and z - axes multiplied by a unit vector (i, j, k) . For such cases, the dot product of two of these vectors $\{a\}$ and $\{b\}$ corresponds to the product of their magnitudes and the cosine of the angle between their tails as in $\{a\} \cdot \{b\} = ab \cos \theta$.

The cross product yields another vector, $\{c\} = \{a\} \times \{b\}$, which is perpendicular to the plane defined by $\{a\}$ and $\{b\}$ such that its direction is specified by the right-hand rule. Develop an M-file function that is passed two such vectors and returns θ , $\{c\}$ and the magnitude of $\{c\}$ and generates a three-dimensional plot of the three vectors $\{a\}$, $\{b\}$ and $\{c\}$ with their origins at zero. Use dashed lines for $\{a\}$ and $\{b\}$ and a solid line for $\{c\}$. Test your function for the following cases:

- (a) $\mathbf{a} = [6 \ 4 \ 2]$; $\mathbf{b} = [2 \ 6 \ 4]$;
- (b) $\mathbf{a} = [3 \ 2 \ -6]$; $\mathbf{b} = [4 \ -3 \ 1]$;
- (c) $\mathbf{a} = [2 \ -2 \ 1]$; $\mathbf{b} = [4 \ 2 \ -4]$;
- (d) $\mathbf{a} = [-1 \ 0 \ 0]$; $\mathbf{b} = [0 \ -1 \ 0]$;