

Equivariant K-stability under finite group action

Ziwen Zhu

Based on joint work with Yuchen Liu

University of Utah

ZAG Seminar, July 30, 2020

Settings

- We will work over the field of complex numbers \mathbb{C} .

Settings

- We will work over the field of complex numbers \mathbb{C} .
- A \mathbb{Q} -Fano variety is a normal projective variety with at worst klt singularities and the anti-canonical divisor \mathbb{Q} -Cartier and ample.

Settings

- We will work over the field of complex numbers \mathbb{C} .
- A \mathbb{Q} -Fano variety is a normal projective variety with at worst klt singularities and the anti-canonical divisor \mathbb{Q} -Cartier and ample.
- A log Fano pair is a pair with at worst klt singularities and the anti-log canonical divisor ample.

Settings

- We will work over the field of complex numbers \mathbb{C} .
- A \mathbb{Q} -Fano variety is a normal projective variety with at worst klt singularities and the anti-canonical divisor \mathbb{Q} -Cartier and ample.
- A log Fano pair is a pair with at worst klt singularities and the anti-log canonical divisor ample.
- The dimension of X is denoted by n .

Motivation

- (Chen-Donaldson-Sun, Tian) A Fano Manifold is Kähler-Einstein if and only if it is K-polystable.

Motivation

- (Chen-Donaldson-Sun, Tian) A Fano Manifold is Kähler-Einstein if and only if it is K-polystable.
- In general it is not easy to verify K-stability.

Motivation

- (Chen-Donaldson-Sun, Tian) A Fano Manifold is Kähler-Einstein if and only if it is K-polystable.
- In general it is not easy to verify K-stability.
- Equivariant K-stability is defined via equivariant test configurations.

Motivation

- (Chen-Donaldson-Sun, Tian) A Fano Manifold is Kähler-Einstein if and only if it is K-polystable.
- In general it is not easy to verify K-stability.
- Equivariant K-stability is defined via equivariant test configurations.
- For \mathbb{Q} -Fano varieties with large symmetry, checking equivariant K-stability is sometimes easier. (E.g. \mathbb{P}^n , V_{22}^*)

Motivation

- (Chen-Donaldson-Sun, Tian) A Fano Manifold is Kähler-Einstein if and only if it is K-polystable.
- In general it is not easy to verify K-stability.
- Equivariant K-stability is defined via equivariant test configurations.
- For \mathbb{Q} -Fano varieties with large symmetry, checking equivariant K-stability is sometimes easier. (E.g. \mathbb{P}^n , V_{22}^*)

Conjecture

Let X be a \mathbb{Q} -Fano variety and G a reductive group action on X . If X is G -equivariantly K-semistable (resp. G -equivariantly K-polystable), then X is K-semistable (resp. K-polystable).

Solution to the equivariant K-stability conjecture:

- (Datar-Székelyhidi) Smooth case;

Solution to the equivariant K-stability conjecture:

- (Datar-Székelyhidi) Smooth case;
- (Li-Xu, Li-Wang-Xu) Torus action;

Solution to the equivariant K-stability conjecture:

- (Datar-Székelyhidi) Smooth case;
- (Li-Xu, Li-Wang-Xu) Torus action;
- (Liu-Z) Finite group action;

Recent progress

Solution to the equivariant K-stability conjecture:

- (Datar-Székelyhidi) Smooth case;
- (Li-Xu, Li-Wang-Xu) Torus action;
- (Liu-Z) Finite group action;
- (Zhuang) General case.

Recent progress

Solution to the equivariant K-stability conjecture:

- (Datar-Székelyhidi) Smooth case;
- (Li-Xu, Li-Wang-Xu) Torus action;
- (Liu-Z) Finite group action;
- (Zhuang) General case.

Equivariant test configuration

A **test configuration** $(\mathcal{X}, \mathcal{L})$ of $(X, -rK_X)$ is a \mathbb{G}_m -equivariant degeneration of X over \mathbb{A}^1 :

$$\begin{array}{ccccc} X \times \{1\} & \hookrightarrow & X \times \mathbb{C}^* & \hookrightarrow & \mathcal{X} \\ \downarrow & & \downarrow & & \downarrow \\ \{1\} & \hookrightarrow & \mathbb{C}^* & \hookrightarrow & \mathbb{A}^1. \end{array}$$

Equivariant test configuration

A **test configuration** $(\mathcal{X}, \mathcal{L})$ of $(X, -rK_X)$ is a \mathbb{G}_m -equivariant degeneration of X over \mathbb{A}^1 :

$$\begin{array}{ccccc} X \times \{1\} & \hookrightarrow & X \times \mathbb{C}^* & \hookrightarrow & \mathcal{X} \\ \downarrow & & \downarrow & & \downarrow \\ \{1\} & \hookrightarrow & \mathbb{C}^* & \hookrightarrow & \mathbb{A}^1. \end{array}$$

There is an induced G -action on $X \times \mathbb{C}^*$:

$$g(x, t) = (\lambda_t \circ g \circ \lambda_t^{-1}(x), t), \quad g \in G.$$

where λ_t is the 1-parameter subgroup inducing the test configuration.

Equivariant test configuration

A **test configuration** $(\mathcal{X}, \mathcal{L})$ of $(X, -rK_X)$ is a \mathbb{G}_m -equivariant degeneration of X over \mathbb{A}^1 :

$$\begin{array}{ccccc} X \times \{1\} & \hookrightarrow & X \times \mathbb{C}^* & \hookrightarrow & \mathcal{X} \\ \downarrow & & \downarrow & & \downarrow \\ \{1\} & \hookrightarrow & \mathbb{C}^* & \hookrightarrow & \mathbb{A}^1. \end{array}$$

There is an induced G -action on $X \times \mathbb{C}^*$:

$$g(x, t) = (\lambda_t \circ g \circ \lambda_t^{-1}(x), t), \quad g \in G.$$

where λ_t is the 1-parameter subgroup inducing the test configuration.

The test configuration is called **G -equivariant** if the above G -action extends to an action on \mathcal{X} commuting with the \mathbb{G}_m -action.

Intersection formula of generalized Futaki invariant

There is a natural $G \times \mathbb{G}_m$ -equivariant compactification $(\overline{\mathcal{X}}, \overline{\mathcal{L}})$ of $(\mathcal{X}, \mathcal{L})$ over \mathbb{P}^1 .

Intersection formula of generalized Futaki invariant

There is a natural $G \times \mathbb{G}_m$ -equivariant compactification $(\overline{\mathcal{X}}, \overline{\mathcal{L}})$ of $(\mathcal{X}, \mathcal{L})$ over \mathbb{P}^1 .

The **generalized Futaki invariant** of a normal $(\mathcal{X}, \mathcal{L})$ can be computed by

$$\text{Fut}(\mathcal{X}, \mathcal{L}) = \frac{1}{(n+1)(-K_X)^n} \left(\frac{n}{r^{n+1}} \overline{\mathcal{L}}^{n+1} + \frac{n+1}{r^n} \overline{\mathcal{L}}^n \cdot K_{\overline{\mathcal{X}}/\mathbb{P}^1} \right)$$

Norm of a test configuration

Let

$$\begin{array}{ccc} & \mathcal{Y} & \\ p \swarrow & & \searrow q \\ X \times \mathbb{P}^1 & \cdots \cdots \cdots \rightarrow & \overline{\mathcal{X}}. \end{array}$$

be any resolution of the graph of $X \times \mathbb{P}^1 \dashrightarrow \overline{\mathcal{X}}$.

Norm of a test configuration

Let

$$\begin{array}{ccc} & \mathcal{Y} & \\ p \swarrow & & \searrow q \\ X \times \mathbb{P}^1 & \cdots \cdots \cdots \rightarrow & \overline{\mathcal{X}}. \end{array}$$

be any resolution of the graph of $X \times \mathbb{P}^1 \dashrightarrow \overline{\mathcal{X}}$.

The **Non-Archimedean norm** of $(\mathcal{X}, \mathcal{L})$ can be computed by

$$J^{\text{NA}}(\mathcal{X}, \mathcal{L}) := \frac{p^*(-K_{X \times \mathbb{P}^1 / \mathbb{P}^1})^n \cdot q^* \bar{\mathcal{L}}}{(-K_X)^n} - \frac{\bar{\mathcal{L}}^{n+1}}{(n+1)(-rK_X)^n}.$$

Norm of a test configuration

Let

$$\begin{array}{ccc} & \mathcal{Y} & \\ p \swarrow & & \searrow q \\ X \times \mathbb{P}^1 & \cdots \cdots \cdots \rightarrow & \overline{\mathcal{X}}. \end{array}$$

be any resolution of the graph of $X \times \mathbb{P}^1 \dashrightarrow \overline{\mathcal{X}}$.

The **Non-Archimedean norm** of $(\mathcal{X}, \mathcal{L})$ can be computed by

$$J^{\text{NA}}(\mathcal{X}, \mathcal{L}) := \frac{p^*(-K_{X \times \mathbb{P}^1 / \mathbb{P}^1})^n \cdot q^* \bar{\mathcal{L}}}{(-K_X)^n} - \frac{\bar{\mathcal{L}}^{n+1}}{(n+1)(-rK_X)^n}.$$

$J^{\text{NA}}(\mathcal{X}, \mathcal{L})$ is one of the several equivalent norms of test configurations that can be used to define uniform K-stability.

Equivariant K-stability

X is called

- **(G -equivariantly) K-semistable** if $\text{Fut}(\mathcal{X}, \mathcal{L}) \geq 0$ for any (G -equivariant) test configuration $(\mathcal{X}, \mathcal{L})$;

Equivariant K-stability

X is called

- **(G -equivariantly) K-semistable** if $\text{Fut}(\mathcal{X}, \mathcal{L}) \geq 0$ for any (G -equivariant) test configuration $(\mathcal{X}, \mathcal{L})$;
- **(G -equivariantly) K-polystable** if $\text{Fut}(\mathcal{X}, \mathcal{L}) \geq 0$ for any (G -equivariant) test configuration $(\mathcal{X}, \mathcal{L})$, and $\text{Fut}(\mathcal{X}, \mathcal{L}) = 0$ if and only if $\mathcal{X} \simeq X \times \mathbb{A}^1$;

Equivariant K-stability

X is called

- **(G -equivariantly) K-semistable** if $\text{Fut}(\mathcal{X}, \mathcal{L}) \geq 0$ for any (G -equivariant) test configuration $(\mathcal{X}, \mathcal{L})$;
- **(G -equivariantly) K-polystable** if $\text{Fut}(\mathcal{X}, \mathcal{L}) \geq 0$ for any (G -equivariant) test configuration $(\mathcal{X}, \mathcal{L})$, and $\text{Fut}(\mathcal{X}, \mathcal{L}) = 0$ if and only if $\mathcal{X} \simeq X \times \mathbb{A}^1$;
- **uniformly (G -equivariantly) K-stable** if there exists $\delta \in (0, 1)$, such that $\text{Fut}(\mathcal{X}, \mathcal{L}) \geq \delta J^{\text{NA}}(\mathcal{X}, \mathcal{L})$ for any (G -equivariant) test configuration $(\mathcal{X}, \mathcal{L})$.

Equivariant K-stability

X is called

- **(G -equivariantly) K-semistable** if $\text{Fut}(\mathcal{X}, \mathcal{L}) \geq 0$ for any (G -equivariant) test configuration $(\mathcal{X}, \mathcal{L})$;
- **(G -equivariantly) K-polystable** if $\text{Fut}(\mathcal{X}, \mathcal{L}) \geq 0$ for any (G -equivariant) test configuration $(\mathcal{X}, \mathcal{L})$, and $\text{Fut}(\mathcal{X}, \mathcal{L}) = 0$ if and only if $\mathcal{X} \simeq X \times \mathbb{A}^1$;
- **uniformly (G -equivariantly) K-stable** if there exists $\delta \in (0, 1)$, such that $\text{Fut}(\mathcal{X}, \mathcal{L}) \geq \delta J^{\text{NA}}(\mathcal{X}, \mathcal{L})$ for any (G -equivariant) test configuration $(\mathcal{X}, \mathcal{L})$.

Note: If we take $G = \{e\}$, then we recover the usual K-stability notions.

Examples (projective space)

- The projective space \mathbb{P}^n is K-polystable, but not uniformly K-stable.

Examples (projective space)

- The projective space \mathbb{P}^n is K-polystable, but not uniformly K-stable.
- Consider $G = \text{Aut}(\mathbb{P}^n) = \text{PGL}(n+1)$.

Examples (projective space)

- The projective space \mathbb{P}^n is K-polystable, but not uniformly K-stable.
- Consider $G = \operatorname{Aut}(\mathbb{P}^n) = \operatorname{PGL}(n+1)$.
 \mathbb{P}^n is uniformly G -equivariantly K-stable, since there is no nontrivial G -equivariant test configuration for \mathbb{P}^n .

Examples (projective space)

- The projective space \mathbb{P}^n is K-polystable, but not uniformly K-stable.
- Consider $G = \operatorname{Aut}(\mathbb{P}^n) = \operatorname{PGL}(n+1)$.
 \mathbb{P}^n is uniformly G -equivariantly K-stable, since there is no nontrivial G -equivariant test configuration for \mathbb{P}^n .
- Therefore the equivariant K-stability conjecture is not true for uniform K-stability.

Examples (Tian's criterion)

For a \mathbb{Q} -Fano variety X with a group action G , we can define its G -equivariant alpha invariant

$$\alpha_G(X) = \inf_m \left\{ \text{lct} \left(X, \frac{1}{m} \Sigma \right) \mid G\text{-invariant } \Sigma \subset | -mK_X | \right\}.$$

Examples (Tian's criterion)

For a \mathbb{Q} -Fano variety X with a group action G , we can define its G -equivariant alpha invariant

$$\alpha_G(X) = \inf_m \left\{ \text{lct} \left(X, \frac{1}{m} \Sigma \right) \mid G\text{-invariant } \Sigma \subset | -mK_X | \right\}.$$

Theorem (Odaka-Sano)

If $\alpha_G(X) \geq$ (resp. $>$) $n/(n+1)$, then X is G -equivariantly K -semistable (uniformly G -equivariantly K -stable).

Examples (Fano threefolds of degree 22)

Example (Cheltsov-Shramov)

V_{22}^* denotes a class of Fano threefolds of degree 22 admitting G -action with $G = \mathbb{C}^* \rtimes \mathbb{Z}/2$. All but two of them have G -equivariant alpha invariants to be $4/5$.

Examples (Fano threefolds of degree 22)

Example (Cheltsov-Shramov)

V_{22}^* denotes a class of Fano threefolds of degree 22 admitting G -action with $G = \mathbb{C}^* \rtimes \mathbb{Z}/2$. All but two of them have G -equivariant alpha invariants to be $4/5$.

According to the equivariant Tian's criterion, those $X \in V_{22}^*$ with $\alpha_G(X) = 4/5$ are uniformly G -equivariantly K-stable and hence K-polystable (Kähler-Einstein) due to the equivariant K-stability conjecture.

Valuative criterion

Let E be a prime divisor over X with a proper birational morphism

$$E \subset Y \xrightarrow{\pi} X.$$

Valuative criterion

Let E be a prime divisor over X with a proper birational morphism

$$E \subset Y \xrightarrow{\pi} X.$$

Denote by $A_X(E)$ the log discrepancy of E , and

$$S_X(E) := \frac{1}{(-K_X)^n} \int_0^{\tau(E)} \text{vol}_Y(\pi^*(-K_X) - xE) \, dx,$$

Valuative criterion

Let E be a prime divisor over X with a proper birational morphism

$$E \subset Y \xrightarrow{\pi} X.$$

Denote by $A_X(E)$ the log discrepancy of E , and

$$S_X(E) := \frac{1}{(-K_X)^n} \int_0^{\tau(E)} \text{vol}_Y(\pi^*(-K_X) - xE) \, dx,$$

where

$$\tau(E) = \sup\{t \mid \text{vol}_Y(\pi^*(-K_X) - tE) > 0\}$$

is the pseudo-effective threshold of E .

Valuative criterion (Cont'd)

The following two invariants are involved in the valuative criterion:

$$\beta(E) = A_X(E) - S_X(E), \quad j(E) = \tau(E) - S_X(E).$$

Valuative criterion (Cont'd)

The following two invariants are involved in the valuative criterion:

$$\beta(E) = A_X(E) - S_X(E), \quad j(E) = \tau(E) - S_X(E).$$

Theorem (Fujita, Li)

A Fano variety X is

- ① *K -semistable if and only if $\beta(E) \geq 0$ for every prime divisor E over X ;*
- ② *uniformly K -stable if and only if there exists $\delta \in (0, 1)$, such that $\beta(E) \geq \delta j(E)$ for every prime divisor E over X .*

Pseudovaluations

Let $G < \text{Aut}(X)$ be a group action on X . For any valuation v on X , we define

$$G \cdot v := \inf_{g \in G} g \cdot v,$$

where $g \cdot v$ is the valuation given by $g \cdot v(f) = v(f \circ g)$ for any $f \in \mathbb{C}(X)$.

Pseudovaluations

Let $G < \text{Aut}(X)$ be a group action on X . For any valuation v on X , we define

$$G \cdot v := \inf_{g \in G} g \cdot v,$$

where $g \cdot v$ is the valuation given by $g \cdot v(f) = v(f \circ g)$ for any $f \in \mathbb{C}(X)$.

For any nonnegative real number x , the ideal sheaf

$$\mathfrak{a}_x(G \cdot v) = \bigcap_{g \in G} \mathfrak{a}_x(g \cdot v)$$

collects regular functions of vanishing order at least x with respect to all $g \cdot v$'s.

Equivariant valuative criterion

Let E be a divisor over X . Define

$$S_X^G(E) := \frac{1}{(-K_X)^n} \int_0^{\tau^G(E)} \text{vol}_X(\mathcal{O}_X(-K_X) \otimes \mathfrak{a}_x(G \cdot \text{ord}_E)) \, dx,$$

where

$$\tau^G(E) := \sup\{t > 0 \mid \text{vol}_X(\mathcal{O}_X(-K_X) \otimes \mathfrak{a}_t(G \cdot \text{ord}_E)) > 0\}.$$

Equivariant valuative criterion

Let E be a divisor over X . Define

$$S_X^G(E) := \frac{1}{(-K_X)^n} \int_0^{\tau^G(E)} \text{vol}_X(\mathcal{O}_X(-K_X) \otimes \mathfrak{a}_x(G \cdot \text{ord}_E)) \, dx,$$

where

$$\tau^G(E) := \sup\{t > 0 \mid \text{vol}_X(\mathcal{O}_X(-K_X) \otimes \mathfrak{a}_t(G \cdot \text{ord}_E)) > 0\}.$$

The equivariant invariants we need for the equivariant valuative criterion are

$$\beta^G(E) = A_X(E) - S_X^G(E), \quad j^G(E) = \tau^G(E) - S_X^G(E).$$

Equivariant valuative criterion

Let E be a divisor over X . Define

$$S_X^G(E) := \frac{1}{(-K_X)^n} \int_0^{\tau^G(E)} \text{vol}_X(\mathcal{O}_X(-K_X) \otimes \mathfrak{a}_x(G \cdot \text{ord}_E)) \, dx,$$

where

$$\tau^G(E) := \sup\{t > 0 \mid \text{vol}_X(\mathcal{O}_X(-K_X) \otimes \mathfrak{a}_t(G \cdot \text{ord}_E)) > 0\}.$$

The equivariant invariants we need for the equivariant valuative criterion are

$$\beta^G(E) = A_X(E) - S_X^G(E), \quad j^G(E) = \tau^G(E) - S_X^G(E).$$

Note that the above invariants coincide with the usual ones if E is G -invariant.

Equivariant valuative criterion (Cont'd)

A prime divisor E over X is **of finite orbit** if the orbit of the valuation ord_E under G -action is finite.

Equivariant valuative criterion (Cont'd)

A prime divisor E over X is **of finite orbit** if the orbit of the valuation ord_E under G -action is finite.

Theorem (Z)

A Fano variety X is

- 1 G -equivariantly K -semistable if and only if $\beta^G(E) \geq 0$ for every finite-orbit prime divisor E over X ;
- 2 uniformly G -equivariantly K -stable if and only if there exists $\delta \in (0, 1)$, such that $\beta^G(E) \geq \delta j^G(E)$ for every finite-orbit prime divisor E over X .

Quotient under finite group action

For the rest of the talk, we assume G is finite.

Quotient under finite group action

For the rest of the talk, we assume G is finite.

Let $\sigma : X \rightarrow Y := X/G$ be the quotient map.

Quotient under finite group action

For the rest of the talk, we assume G is finite.

Let $\sigma : X \rightarrow Y := X/G$ be the quotient map. We have the following relation of canonical divisors

$$K_X = \sigma^* K_Y + \sum (e_R - 1)R,$$

where R runs through all ramification divisors and e_R is the ramification index of R .

Quotient under finite group action

For the rest of the talk, we assume G is finite.

Let $\sigma : X \rightarrow Y := X/G$ be the quotient map. We have the following relation of canonical divisors

$$K_X = \sigma^* K_Y + \sum (e_R - 1)R,$$

where R runs through all ramification divisors and e_R is the ramification index of R . For each branch divisor B_i on Y , denote by e_i the ramification index of R with $\sigma(R) = B_i$. Then we have

$$K_X = \sigma^*(K_Y + B),$$

where $B = \sum_i \left(1 - \frac{1}{e_i}\right) B_i$.

K-stability of quotient

Following the notation in the previous slide, (Y, B) is a log Fano pair if X is Fano. All previous results work for log Fano pairs by replacing canonical divisors with log canonical divisors.

K-stability of quotient

Following the notation in the previous slide, (Y, B) is a log Fano pair if X is Fano. All previous results work for log Fano pairs by replacing canonical divisors with log canonical divisors.

Proposition (Liu-Z)

Under the above notation, X is G -equivariantly K -semistable (resp. uniformly G -equivariantly K -stable) if and only if (Y, B) is K -semistable (resp. uniformly K -stable).

K-stability of quotient

Following the notation in the previous slide, (Y, B) is a log Fano pair if X is Fano. All previous results work for log Fano pairs by replacing canonical divisors with log canonical divisors.

Proposition (Liu-Z)

Under the above notation, X is G -equivariantly K -semistable (resp. uniformly G -equivariantly K -stable) if and only if (Y, B) is K -semistable (resp. uniformly K -stable).

Idea of the proof:

- "Only if" part: comparing beta invariants of corresponding prime divisors;
- "If" part: comparing generalized Futaki invariants of corresponding test configurations.

Main Result

Theorem (Liu-Z)

Let X be a \mathbb{Q} -Fano variety and G a finite group action on X . If X is G -equivariantly K -semistable (resp. G -equivariantly K -polystable), then X is K -semistable (resp. K -polystable).

Main Result

Theorem (Liu-Z)

Let X be a \mathbb{Q} -Fano variety and G a finite group action on X . If X is G -equivariantly K -semistable (resp. G -equivariantly K -polystable), then X is K -semistable (resp. K -polystable).

Idea of the proof (K -semistable case):

Main Result

Theorem (Liu-Z)

Let X be a \mathbb{Q} -Fano variety and G a finite group action on X . If X is G -equivariantly K -semistable (resp. G -equivariantly K -polystable), then X is K -semistable (resp. K -polystable).

Idea of the proof (K -semistable case):

X is G -equivariantly K -semistable $\Rightarrow (Y, B)$ is K -semistable

Main Result

Theorem (Liu-Z)

Let X be a \mathbb{Q} -Fano variety and G a finite group action on X . If X is G -equivariantly K -semistable (resp. G -equivariantly K -polystable), then X is K -semistable (resp. K -polystable).

Idea of the proof (K -semistable case):

X is G -equivariantly K -semistable $\Rightarrow (Y, B)$ is K -semistable

$\xrightarrow{\text{Blum-Liu}} (Y, B + \epsilon D')$ is uniformly K -stable, for some

$D' \in |-(K_Y + B)|_{\mathbb{Q}}$

Main Result

Theorem (Liu-Z)

Let X be a \mathbb{Q} -Fano variety and G a finite group action on X . If X is G -equivariantly K -semistable (resp. G -equivariantly K -polystable), then X is K -semistable (resp. K -polystable).

Idea of the proof (K -semistable case):

X is G -equivariantly K -semistable $\Rightarrow (Y, B)$ is K -semistable

$\xrightarrow{\text{Blum-Liu}} (Y, B + \epsilon D')$ is uniformly K -stable, for some

$D' \in |-(K_Y + B)|_{\mathbb{Q}}$

$\xrightarrow{\text{Uniform YTD (Li-Tian-Wang)}} (Y, B + \epsilon D')$ is Kähler-Einstein

Main Result

Theorem (Liu-Z)

Let X be a \mathbb{Q} -Fano variety and G a finite group action on X . If X is G -equivariantly K -semistable (resp. G -equivariantly K -polystable), then X is K -semistable (resp. K -polystable).

Idea of the proof (K -semistable case):

X is G -equivariantly K -semistable $\Rightarrow (Y, B)$ is K -semistable

$\xrightarrow{\text{Blum-Liu}} (Y, B + \epsilon D')$ is uniformly K -stable, for some

$D' \in |-(K_Y + B)|_{\mathbb{Q}}$

$\xrightarrow{\text{Uniform YTD (Li-Tian-Wang)}} (Y, B + \epsilon D')$ is Kähler-Einstein

$\Rightarrow (X, \epsilon D)$ is Kähler-Einstein, where $D = \sigma^* D' \in |-K_X|_{\mathbb{Q}}$

Main Result

Theorem (Liu-Z)

Let X be a \mathbb{Q} -Fano variety and G a finite group action on X . If X is G -equivariantly K -semistable (resp. G -equivariantly K -polystable), then X is K -semistable (resp. K -polystable).

Idea of the proof (K -semistable case):

X is G -equivariantly K -semistable $\Rightarrow (Y, B)$ is K -semistable

$\xrightarrow{\text{Blum-Liu}} (Y, B + \epsilon D')$ is uniformly K -stable, for some

$D' \in |-(K_Y + B)|_{\mathbb{Q}}$

$\xrightarrow{\text{Uniform YTD (Li-Tian-Wang)}} (Y, B + \epsilon D')$ is Kähler-Einstein

$\Rightarrow (X, \epsilon D)$ is Kähler-Einstein, where $D = \sigma^* D' \in |-K_X|_{\mathbb{Q}}$

$\xrightarrow{\text{Berman}} (X, \epsilon D)$ is K -semistable $\xrightarrow{\text{Blum-Liu}} X$ is K -semistable.

Main Result

Theorem (Liu-Z)

Let X be a \mathbb{Q} -Fano variety and G a finite group action on X . If X is G -equivariantly K -semistable (resp. G -equivariantly K -polystable), then X is K -semistable (resp. K -polystable).

Idea of the proof (K -semistable case):

X is G -equivariantly K -semistable $\Rightarrow (Y, B)$ is K -semistable

$\xrightarrow{\text{Blum-Liu}} (Y, B + \epsilon D')$ is uniformly K -stable, for some

$D' \in |-(K_Y + B)|_{\mathbb{Q}}$

$\xrightarrow{\text{Uniform YTD (Li-Tian-Wang)}} (Y, B + \epsilon D')$ is Kähler-Einstein

$\Rightarrow (X, \epsilon D)$ is Kähler-Einstein, where $D = \sigma^* D' \in |-K_X|_{\mathbb{Q}}$

$\xrightarrow{\text{Berman}} (X, \epsilon D)$ is K -semistable $\xrightarrow{\text{Blum-Liu}} X$ is K -semistable.

Note: The proof of the uniform YTD conjecture in L-T-W is analytic.

Idea of Proof (K-polystable case)

Assume X is strictly K-semistable, then there is a special test configuration \mathcal{X} degenerating X to some X_0 which is K-polystable.

Idea of Proof (K-polystable case)

Assume X is strictly K-semistable, then there is a special test configuration \mathcal{X} degenerating X to some X_0 which is K-polystable.

Consider an étale neighborhood of $[X_0]$ in the corresponding K-moduli stack of the form $[\mathrm{Spec} A/H]$ where $H = \mathrm{Aut}(X_0)$ and $G < \mathrm{Aut}(X) < H$.

A test configuration \mathcal{X} degenerating X to X_0 corresponds to a morphism $\phi : [\mathbb{A}^1/\mathbb{G}_m] \rightarrow [\mathrm{Spec} A/H]$ with $\phi(1) = [X]$ and $\phi(0) = [X_0]$.

Idea of Proof (K-polystable case)

Assume X is strictly K-semistable, then there is a special test configuration \mathcal{X} degenerating X to some X_0 which is K-polystable.

Consider an étale neighborhood of $[X_0]$ in the corresponding K-moduli stack of the form $[\mathrm{Spec} A/H]$ where $H = \mathrm{Aut}(X_0)$ and $G < \mathrm{Aut}(X) < H$.

A test configuration \mathcal{X} degenerating X to X_0 corresponds to a morphism $\phi : [\mathbb{A}^1/\mathbb{G}_m] \rightarrow [\mathrm{Spec} A/H]$ with $\phi(1) = [X]$ and $\phi(0) = [X_0]$.

Equivalently, we have a 1-parameter subgroup: $\lambda : \mathbb{G}_m \rightarrow H$ with

$$\lim_{t \rightarrow 0} \lambda(t) \cdot [X] = [X_0].$$

Idea of Proof (K-polystable case)

Our goal is to find a G -equivariant ϕ' such that we get a G -equivariant test configuration \mathcal{X}' with $\text{Fut}(\mathcal{X}') = 0$.

Idea of Proof (K-polystable case)

Our goal is to find a G -equivariant ϕ' such that we get a G -equivariant test configuration \mathcal{X}' with $\text{Fut}(\mathcal{X}') = 0$.

A result of Luna about closedness of orbits \Rightarrow we have a 1-parameter subgroup: $\lambda' : \mathbb{G}_m \rightarrow Z_H(G)$ with

$$\lim_{t \rightarrow 0} \lambda'(t) \cdot [X] = [X_0],$$

where $Z_H(G)$ is the centralizer of G inside H .

Idea of Proof (K-polystable case)

Our goal is to find a G -equivariant ϕ' such that we get a G -equivariant test configuration \mathcal{X}' with $\text{Fut}(\mathcal{X}') = 0$.

A result of Luna about closedness of orbits \Rightarrow we have a 1-parameter subgroup: $\lambda' : \mathbb{G}_m \rightarrow Z_H(G)$ with

$$\lim_{t \rightarrow 0} \lambda'(t) \cdot [X] = [X_0],$$

where $Z_H(G)$ is the centralizer of G inside H .

This gives us the desired G -equivariant test configuration \mathcal{X}' .

Idea of Proof (K-polystable case)

Our goal is to find a G -equivariant ϕ' such that we get a G -equivariant test configuration \mathcal{X}' with $\text{Fut}(\mathcal{X}') = 0$.

A result of Luna about closedness of orbits \Rightarrow we have a 1-parameter subgroup: $\lambda' : \mathbb{G}_m \rightarrow Z_H(G)$ with

$$\lim_{t \rightarrow 0} \lambda'(t) \cdot [X] = [X_0],$$

where $Z_H(G)$ is the centralizer of G inside H .

This gives us the desired G -equivariant test configuration \mathcal{X}' .

\Rightarrow Contradiction, and hence X is K-polystable.

Thank you!