
- · Contitewing + stability
- sensitively to perturbed data
- o ex: scaler accitour.

Control of the second


Part III: Conditions + Stability:

A problem is well-conditioned if small changes in the data led to


rba el n agando lloma



begin about volve



An algorithm or method is called stable if it always produces a solly to a number produce.



of a proble os ill-conditioned, stability of the absorption is not useful.

Protoun conditioning is essential to nomorical methods byc we work with rande date.

If an algorithm is applied to a protein with data de to opt a sola so that sisce).

De or introduce in the relative error 11 sco) - sca)11 when a sounded (or otherwise pollutes) data.

Ex 1: Scales addition

· data: and ER

& soution: S = ax b

· pertonal deta: ~= a+ Aa, ~= b+Ab

· performed soly: \$= 2+6

relative unon: $\frac{|8-3|}{|3|} = \frac{|\tilde{a}+\tilde{b}|}{|3|} = \frac{|\Delta a+\Delta b|}{|a+b|} = \frac{|\Delta a|+|\Delta b|}{|a+b|} = \frac{1}{|a+b|} \left(|a| \frac{|\Delta a|}{|a|} + |b| \frac{|\Delta b|}{|b|} \right)$

ed 18-81 L lat-161 > max { [M], (Ab) }

181 (101)

181 (101)

182 (101)

182 (101)

182 (101)

183 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184 (101)

184

C: lal 1/5/ magrifua the poterbation to the data.

if sign(a) = sign(b), c=1

if I a+bl « Ial + Ibl Ccanellation vior), c>>1: c "blown op" pertubation of deta.

=D subtractor with cancellation is ill-conditioned!

More: precises (book rotet un)

Let T(x) be a small potarboth that of a $T(f) = f(x+T\alpha) - f(x) \qquad ("f" is soon "a" is down)$

The absolute and from nowber $\hat{K} = \lim_{n \to \infty} \frac{s_n}{n + n} \frac{1181}{n + n}$

It losts like a southern arivolve.

JR 784 563) = 5 the Southon of f at M: 5is = Tom;

then TF = 5635 or All and & = 1156311.

Usocho we are conwined with the teletire condition maker

15: 6(x) = lim sup (15415) (1541) / 1541)

charges magnified at most by

CAMPHON N