

ETAMv2.0: Energetic–Temporal–Environmental Activity Model, Version 2.0

Shao Kun <11947910734@qq.com>

2026-02-22

1 Notation and Conventions

All symbols are defined as follows unless otherwise noted:

- **A–T–E** (Spatial-Temporal-Energetic): primitive ontological basis **A** for *spatial-architectural activity*, **T** for *temporal-slot activation*, **E** for *energo-material response*
- **XR** (ATE Basis Encoding): discrete coordinate representation of the **A–T–E** (Spatial-Temporal-Energetic) basis; $R = [\mathbf{n}_1, \dots, \mathbf{n}_N]^\top \in \mathbb{Z}^{N \times 3}$
- **icR** (Complex Generator: $f(R) = x(R) + i c(R)$): complex-valued generator $f(R) = x(R) + i c(R)$, where x is the *covariant observable*, c the *contravariant lifting potential*

2 Introduction

The **Energetic–Temporal–Environmental Activity Model (ETAM)** is a physics-informed, differentiable framework for predicting macroscopic scalar outputs Y from discrete-continuous hybrid inputs. Version 2.0 introduces unified slot-based encoding, complex-valued response functions, and adaptive multilevel hierarchy driven by the imaginary component.

Clarification of novelty: The **A–T–E** (Spatial-Temporal-Energetic) triad is not a re-labeling of conventional physical quantities. It is a *primitive ontological basis* defined by its role in the **icR** (Complex Generator: $f(R) = x(R) + i c(R)$) generator: **A** encodes spatial-architectural constraints, **T** encodes temporal-slot resolution, **E** encodes energo-material coupling. No existing model uses this triad as a unified coordinate system.

This aligns with Rosen’s relational ontology: **A–T–E** (Spatial-Temporal-Energetic) does not describe *what things are*, but *how they co-constitute observables through constrained coordination* ?.

3 Core Output Structure

The model defines a dual-domain mapping:

$$Y = A \cdot T \cdot E \quad (1)$$

$$f(R) = x(R) + i c(R) \quad (2)$$

where $Y \in \mathbb{R}_{\geq 0}$ is the physical observable (e.g., binding energy in eV), and $f(R) \in \mathbb{C}$ is the **icR** (Complex Generator: $f(R) = x(R) + i c(R)$) generator. Alignment requires $\Re[f(R)] \approx Y$, while $\Im[f(R)] = c(R) \geq 0$ governs scale-level selection.

4 Input Representation R

Each active unit j (atom or functional group) is encoded as a discrete triple:

$$\mathbf{n}_j = (n_{A,j}, n_{T,j}, n_{E,j}) \in \mathbb{Z}_{\geq 0}^3. \quad (3)$$

For N units, the input tensor is:

$$R = [\mathbf{n}_1, \mathbf{n}_2, \dots, \mathbf{n}_N]^\top \in \mathbb{Z}^{N \times 3}. \quad (4)$$

R is not a feature vector it is the *coordinate representation of the A-T-E (Spatial-Temporal-Energetic) basis in discrete state space*. Each row $\mathbf{n}_j$ is a point in the A-T-E (Spatial-Temporal-Energetic) lattice, where $n_{A,j}, n_{T,j}, n_{E,j}$ are *mutually constraining coordinates*, not independent scalars.

5 Component Definitions

5.1 A-value: Spatial-Architectural Activity

Let $\omega(p) \in \{0, 1, 2\}$ denote occupancy (empty/single electron/lone pair) at orbital slot p . Define slot weights $k(p)$:

$$k(p) = \begin{cases} 1 & \text{for } s\text{-orbitals,} \\ 2 & \text{for } p\text{-orbitals,} \\ 3 & \text{for } d\text{-orbitals,} \\ 4 & \text{for } f\text{-orbitals.} \end{cases} \quad (5)$$

The activity constant for unit j is:

$$C_{A,j} = \sum_{p \in \{s,p,d,f\}} \omega(p) k(p). \quad (6)$$

Then:

$$A_j = \alpha_A \cdot C_{A,j}, \quad A = \frac{1}{N} \sum_{j=1}^N A_j, \quad (7)$$

with scaling factor $\alpha_A > 0$.

5.2 T-value: Temporal-Slot Activation

Six discrete time slots $k = 0, \dots, 5$ cover 1×10^{-15} s to 1×10^6 s. Let $\nu(k) \in \{0, 1, 2\}$ indicate activation level. Then:

$$C_{T,j} = \sum_{k=0}^5 \nu(k) w(k), \quad T_j = \alpha_T \cdot C_{T,j}, \quad T = \frac{1}{N} \sum_{j=1}^N T_j. \quad (8)$$

5.3 E-value: Energo-Material Response Field

Four environmental response slots: ionization (e_I), affinity (e_A), polarization (e_P), solvation (e_S). Let $\mu(e) \in \{0, 1, 2\}$. Define:

$$C_{E,j} = \sum_{e \in \{I,A,P,S\}} \mu(e) w_e. \quad (9)$$

Environment-modulated field:

$$\Phi_I(\Xi) = \frac{F^2}{IE}, \quad (10)$$

$$\Phi_P(\Xi) = \frac{\epsilon - 1}{\epsilon + 2} F^2, \quad (11)$$

$$\Phi_S(\Xi) = \frac{\Delta G_{\text{solv}}}{RT}, \quad (12)$$

and:

$$E_j = \alpha_E \cdot C_{E,j} \cdot \left[1 + \sum_e \underbrace{\frac{\mu(e) w_e}{C_{E,j}}}_{\beta_e(C_{E,j})} \cdot \Phi_e(\Xi) \right], \quad E = \frac{1}{N} \sum_{j=1}^N E_j. \quad (13)$$

6 Response Function $f(R)$

The **icR** (Complex Generator: $f(R) = x(R) + i c(R)$) framework ($f(R) = x(R) + i c(R)$) is not an engineering choice for numerical stability. It is a *structural necessity*: only complex-valued representation permits simultaneous encoding of covariant observables (x) and contravariant scale selectors (c) within a single differentiable map.

The icR mapping realizes a *Connes-type spectral geometry*?: discrete **A–T–E** (Spatial-Temporal-Energetic) coordinates R define a finite spectral triple, whose complex extension $f(R)$ yields a differentiable continuum limit.

6.1 Real Part (Covariant Observable)

$$x(R) = \alpha_x \cdot \langle C_A \rangle \cdot \langle C_T \rangle \cdot \langle C_E \rangle, \quad (14)$$

or alternatively via MLP:

$$x(R) = \text{MLP}_{\theta_x}(\log(C_A + \varepsilon), \log(C_T + \varepsilon), \log(C_E + \varepsilon)). \quad (15)$$

6.2 Imaginary Part (Contravariant Lifting Potential)

$$c(R) = \lambda \cdot \left[\frac{\text{Var}(C_A)}{\langle C_A \rangle^2} + \frac{\text{Var}(C_T)}{\langle C_T \rangle^2} \right] \cdot (1 + \eta \cdot S_{\text{env}}), \quad (16)$$

where $S_{\text{env}} = \sum_e \Phi_e(\Xi)$. The non-negative imaginary part $c(R)$ implements Penrose’s ‘ladder of infinity’ on the complex plane?: each threshold τ_k marks a rung where the system’s effective description ascends to a new ontological level.

- $c(R) < \tau_1$: Level 1 (molecular determinism)
- $\tau_1 \leq c(R) < \tau_2$: Level 2 (mesoscale clusters)
- $c(R) \geq \tau_2$: Level 3 (condensed-phase statistical fields)

7 Training Objective

$$\mathcal{L} = \|\Re[f(R)] - Y\|^2 + \gamma \cdot \mathcal{L}_{\text{reg}}(c(R)), \quad (17)$$

where $\mathcal{L}_{\text{reg}}$ enforces level consistency (e.g., entropy penalty on $c(R)$) or other domain-informed constraints.

8 Constrained Inverse Queries

The model supports *constrained inverse queries*: given a partial configuration $\tilde{R}$ and target scalar y_{target} , the set of completions R_{miss} satisfying both energy fidelity and level consistency is non-empty and computable under standard differentiable optimization.

This inverse capability is not an add-on feature. It is a *logical consequence* of the **A–T–E** (Spatial-Temporal-Energetic)–**icR** (Complex Generator: $f(R) = x(R) + i c(R)$) duality: if f is bijective over its domain (up to level constraints), then generation is the algebraic inverse of prediction?.

9 Version Identification

ETAMv2.0 is formally defined by this specification (v2.0, 2026-02-22). Key normative innovations:

1. Unified discrete slot encoding across **A**, **T**, and **E** as a primitive ontological basis;
2. Complex-valued separation of covariant observable (x) and contravariant lifting potential (c);
3. Imaginary part-driven adaptive multilevel hierarchy grounded in Penrose’s ladder;
4. Verifiable scientific discovery via Wigner-style bidirectional invertibility.