

a
Three separate analyses and their objects (schematic)
1 · Inputs and modules
2 · Derived objects
3 · Separate analyses
HF and HCC networks
177 failing LV · 148 tumours
All named modules
23 HF; 20 HCC
Module preservation
both directions; Zsummary
Networks + 15 Tier A sets
→ HF purple; HCC magenta
TCS and module score
mean z; kME-weighted
Score–trait association
TF correlation; survival
15 Tier A sets
explicit membership
ssGSEA per two-arm cohort
pathway scores per sample
Hedges g → mirror
opp. sign; |g| ≥ 0.20; CI≠ 0
b
Module inventory (β = 14)
turquoise
blue
brown
yellow
green
red
black
pink
magenta
purple
greenyellow
tan
salmon
cyan
midnightblue
lightcyan
lightgreen
grey60
lightyellow
royalblue
darkred
darkgreen
darkturquoise
purple (153)
HF
magenta (146)
HCC
0
2,000
4,000
Genes in module
c
Enrichment by module size
HF
HCC
purple
magenta
Tier A fold enrichment
9
6
3
0
100
300
1000
3000
Genes in module (log scale)
d
Module preservation, both directions
dotted lines: Z = 2 and Z = 10
HCC network tested in HF
HF network tested in HCC
Median rank
20
15
10
5
0
magenta (146)
purple (153)
0
5
10
15
Zsummary preservation
e
Preservation classes of named modules, by direction
counts are of named modules; the unassigned grey bin is excluded
HF network tested in HCC
HCC network tested in HF
11
9
3
6
12
2
0%
25%
50%
75%
100%
Fraction of named modules
Preservation class
strong (>10)
weak-moderate (2-10)
none (<2)