

# Frontiers in Differential Geometry, Partial Differential Equations and Mathematical Physics

In Memory of Gu Chaohao

Edited by

**Molin Ge**

Chern Institute of Mathematics, China

**Jiaxing Hong**

Fudan University, China

**Tatsien Li**

Fudan University, China

**Weiping Zhang**

Chern Institute of Mathematics, China

 **World Scientific**

NEW JERSEY • LONDON • SINGAPORE • BEIJING • SHANGHAI • HONG KONG • TAIPEI • CHENNAI

*Published by*

World Scientific Publishing Co. Pte. Ltd.

5 Toh Tuck Link, Singapore 596224

USA office: 27 Warren Street, Suite 401-402, Hackensack, NJ 07601

UK office: 57 Shelton Street, Covent Garden, London WC2H 9HE

### **Library of Congress Cataloging-in-Publication Data**

Frontiers in differential geometry, partial differential equations, and mathematical physics : in memory of Gu Chaozhao / edited by Molin Ge (Chern Institute of Mathematics, China), Jiaxing Hong (Fudan University, China), Tatsien Li (Fudan University, China), & Weiping Zhang (Chern Institute of Mathematics, China).

pages cm

Includes bibliographical references.

ISBN 978-9814578073 (hardcover : alk. paper)

1. Gu, Chaozhao. 2. Geometry, Differential. 3. Differential equations, Partial. 4. Mathematical physics. I. Ge, M. L. (Mo-Lin) II. Hong, Jiaxing, 1942– III. Li, Daqian. IV. Zhang, Weiping, 1964–

QA641.F77 2014

516.3'6--dc23

2013048729

### **British Library Cataloguing-in-Publication Data**

A catalogue record for this book is available from the British Library.

Copyright © 2014 by World Scientific Publishing Co. Pte. Ltd.

*All rights reserved. This book, or parts thereof, may not be reproduced in any form or by any means, electronic or mechanical, including photocopying, recording or any information storage and retrieval system now known or to be invented, without written permission from the publisher.*

For photocopying of material in this volume, please pay a copying fee through the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923, USA. In this case permission to photocopy is not required from the publisher.

Printed in Singapore by World Scientific Printers.

## Contents

<i>Preface</i>	v
A Profile of the Late Professor Gu Chaohao <i>Tatsien Li</i>	1
List of Publications of Gu Chaohao	17
In Memory of Professor Gu Chaohao <i>Xiaqi Ding</i>	27
In Memory of Professor Gu Chaohao <i>Gongqing Zhang (Kung Ching Chang)</i>	29
Stability of E-H Mach Configuration in Pseudo-Steady Compressible Flow <i>Shuxing Chen</i>	35
Incompressible Viscous Fluid Flows with Slip Boundary Conditions and Their Numerical Simulations <i>Ben-yu Guo</i>	49
Global Existence and Uniqueness of the Solution for the Generalized Schrödinger-KdV System <i>Boling Guo, Bolin Ma, Jingjun Zhang</i>	69
Anomaly Cancellation and Modularity <i>Fei Han, Kefeng Liu, Weiping Zhang</i>	87

On Interior Estimates for Mean Curvature of Convex Surfaces in $R^3$ and Its Applications <i>Jiaring Hong</i>	105
Geometric Invariant Theory of the Space — A Modern Approach to Solid Geometry <i>Wu-yi Hsiang</i>	125
Optimal Convergence Rate of the Binomial Tree Scheme for American Options and Their Free Boundaries <i>Lishang Jiang, Jin Liang</i>	153
Rademacher $\Phi$ Function, Jacobi Symbols, Quantum and Classical Invariants of Lens Spaces <i>Bang-He Li, Tian-Jun Li</i>	169
Historical Review on the Roles of Mathematics in the Study of Aerodynamics <i>Jiachun Li</i>	189
Toward Chern–Simons Theory of Complexes on Calabi–Yau Threefolds <i>Jun Li</i>	199
Exact Boundary Synchronization for a Coupled System of Wave Equations <i>Tatsien Li</i>	219
Scaling Limit for Compressible Viscoelastic Fluids <i>Xianpeng Hu, Fanghua Lin</i>	243
Uniqueness Modulo Reduction of Bergman Meromorphic Compactifications of Canonically Embeddable Bergman Manifolds <i>Ngaiming Mok</i>	271
The Application of Conditional Nonlinear Optimal Perturbation to Targeted Observations for Tropical Cyclone Prediction <i>Mu Mu, Feifan Zhou, Xiaohao Qin and Boyu Chen</i>	291
Isometric Immersions in Minkowski Spaces <i>Yi-Bing Shen</i>	327

Remarks on Volume Growth for Minimal Graphs in Higher Codimension <i>Yuanlong Xin</i>	345
Separation of Variables for the Lax Pair of the Bogomolny Equation in 2+1 Dimensional Anti-de Sitter Space-Time <i>Zi-Xiang Zhou</i>	351

# Separation of Variables for the Lax Pair of the Bogomolny Equation in 2+1 Dimensional Anti-de Sitter Space-time

Zi-Xiang Zhou\*

*School of Mathematical Sciences, Fudan University,  
Shanghai 200433, China  
zzzhou@fudan.edu.cn*

A Lax system separating all the variables is reduced from the Lax pair of the Bogomolny equation for the Yang–Mills–Higgs field in 2+1 dimensional anti-de Sitter space-time. It also contains the Bogomolny equation in 2+1 dimensional Minkowski space-time (Ward equation) under further reduction or limit. The Darboux transformation can be constructed in the usual way to get explicit solutions of the Bogomolny equation.

## 1. Introduction

The Bogomolny equation for the Yang–Mills–Higgs field in the 2+1 dimensional anti-de Sitter space-time and Minkowski space-time are known to be integrable.<sup>8,15</sup> They have been studied and solved explicitly by several ways and various kinds of solutions have been obtained<sup>5,8–10,14,15,17,18</sup>

On the other hand, a lot of integrable systems in 2+1 dimensions without spectral parameters can be reduced by separating variables in the Lax pair. A typical way is the nonlinear constraint method,<sup>1,2,11</sup> in which a 2+1 dimensional integrable system can be reduced to several 1+1 dimensional systems.<sup>7</sup> Under this kind of constraint, we can obtain explicit solutions like the dromions<sup>19</sup> and the quasi-periodic solutions<sup>1,20</sup> of the 2+1 dimensional integrable system,

In this paper, we will separate the variables for the Lax pair (with a spectral parameter) of the Bogomolny equation in 2+1 dimensional anti-de

---

\*This work was supported by the National Natural Science Foundation of China (11171073) and the Key Laboratory of Mathematics for Nonlinear Sciences of Ministry of Education of China.

Sitter space-time. The derived system is written in 1+1 dimensional form. The Darboux transformation can be applied to this derived system.

By various limit or reduction process, the results are also valid for the Bogomolny equation in 2+1 dimensional Minkowski space-time<sup>3,5,8</sup> (i.e. Ward equation, or the 2+1 dimensional principal chiral field with Wess–Zumino–Witten term<sup>15</sup>) and the standard 1+1 dimensional principal chiral field.<sup>6,12,13,16</sup>

## 2. Bogomolny Equation in 2+1 Dimensional Anti-de Sitter Space-time

The 2+1 dimensional anti-de Sitter space-time (AdS) of constant curvature  $-1/\rho^2$  ( $\rho > 0$ ) is the hyperboloid

$$U^2 + V^2 - X^2 - Y^2 = \rho^2 \quad (1)$$

in  $\mathbf{R}^{2,2}$  with the metric

$$ds^2 = -dU^2 - dV^2 + dX^2 + dY^2. \quad (2)$$

By defining

$$r = \frac{\rho}{U+X} - \rho + 1, \quad y = \frac{Y}{U+X}, \quad t = -\frac{V}{U+X}, \quad (3)$$

a part of the AdS with  $U+X > 0$  is represented by the Poincaré coordinates  $(r, y, t)$  with  $r > -\rho + 1$  and the metric is

$$\begin{aligned} ds^2 &= \frac{\rho^2}{(r + \rho - 1)^2} (-dt^2 + dr^2 + dy^2) \\ &= \frac{\rho^2}{(x_2 + \rho - 1)^2} (dx_2^2 - 4 dx_1 dx_3) \end{aligned} \quad (4)$$

where  $x_1 = (y + t)/2$ ,  $x_2 = r$ ,  $x_3 = (t - y)/2$ .

For simplicity, denote

$$\gamma = \frac{x_2 + \rho - 1}{\rho}. \quad (5)$$

Here are some basic notions for the Yang–Mills–Higgs field.

Let  $G$  be an  $N \times N$  matrix Lie group,  $\mathcal{G}$  be its Lie algebra. The Yang–Mills–Higgs field in AdS is given by the Yang–Mills potentials  $\{A_\mu\}$  and the Higgs field  $\Phi$  with  $A_\mu \in \mathcal{G}$ ,  $\Phi \in \mathcal{G}$ . Denote  $\partial_\mu = \partial/\partial x_\mu$  where  $\mu = 1, 2, 3$ . The covariant derivative of an  $N$ -dimensional vector  $\psi$  on which  $\mathcal{G}$  acts is  $D_\mu \psi = \partial_\mu \psi + A_\mu \psi$ . The strength of the Yang–Mills field is defined by

$F_{\mu\nu} = [D_\mu, D_\nu] = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu]$ , and the covariant derivative of the Higgs field  $\Phi$  is defined by  $D_\mu \Phi = \partial_\mu \Phi + [A_\mu, \Phi]$ .

For  $g(x) \in G$ , the transformation

$$A'_\mu = gA_\mu g^{-1} - (\partial_\mu g)g^{-1}, \quad \Phi' = g\Phi g^{-1} \quad (6)$$

is called a gauge transformation. Under the gauge transformation, the strength is changed to

$$F'_{\mu\nu} = gF_{\mu\nu}g^{-1}. \quad (7)$$

The Bogomolny equation with the metric (4) is<sup>18</sup>

$$D_1 \Phi = \gamma F_{12}, \quad D_2 \Phi = \frac{\gamma}{2} F_{13}, \quad D_3 \Phi = \gamma F_{23}. \quad (8)$$

By (6) and (7), a gauge transformation transforms a solution of (8) to another solution of it.

**Lemma 1.** *For any Yang-Mills-Higgs field  $(A_\mu, \Phi)$  satisfying the Bogomolny equation (8) on AdS, there exists a gauge transformation such that  $A'_1 = 0$ ,  $A'_2 = \gamma^{-1}\Phi'$  where  $\gamma$  is defined by (5).*

**Proof:** After the gauge transformation given by  $g \in G$ ,

$$\begin{aligned} A'_1 &= gA_1 g^{-1} - (\partial_1 g)g^{-1}, \\ A'_2 - \gamma^{-1}\Phi' &= g(A_2 - \gamma^{-1}\Phi)g^{-1} - (\partial_2 g)g^{-1}. \end{aligned} \quad (9)$$

If we want  $A'_1 = 0$ ,  $A'_2 = \gamma^{-1}\Phi'$ , then  $g$  should satisfy

$$\partial_1 g = gA_1, \quad \partial_2 g = g(A_2 - \gamma^{-1}\Phi). \quad (10)$$

The integrability condition of (10) is

$$\partial_2 A_1 - \partial_1 (A_2 - \gamma^{-1}\Phi) - [A_1, A_2 - \gamma^{-1}\Phi] = 0, \quad (11)$$

which is the first equation of (8). Hence  $g$  always exists. Moreover, since  $A_1 \in \mathcal{G}$  and  $A_2 - \gamma^{-1}\Phi \in \mathcal{G}$ , we can want  $g \in G$ . The lemma is proved.

Thus, we shall always choose the gauge so that  $A_1 = 0$ ,  $A_2 = \gamma^{-1}\Phi$  hold identically.

It was known that (8) has a Lax pair<sup>18</sup>

$$\begin{aligned} &((x_2 + \rho - 1)D_2 + \rho\Phi - (\rho\lambda - 2x_1)D_1)\eta = 0, \\ &\left(D_3 - \frac{\rho\lambda - 2x_1}{x_2 + \rho - 1}D_2 + \frac{\rho(\rho\lambda - 2x_1)}{(x_2 + \rho - 1)^2}\Phi\right)\eta = 0 \end{aligned} \quad (12)$$

which was presented by Ref. 15. This means that (8) is the integrability condition of (12).



Considering the gauge fixed in Lemma 1, (12) can be written as

$$\begin{aligned}\partial_2 \eta &= \sigma(\lambda) \partial_1 \eta - P \eta, \\ \partial_3 \eta &= \sigma(\lambda) \partial_2 \eta - Q \eta\end{aligned}\quad (13)$$

where

$$\sigma(\lambda) = \frac{\rho\lambda - 2x_1}{x_2 + \rho - 1}, \quad (14)$$

$$P = 2\gamma^{-1}\Phi = 2A_2, \quad Q = A_3. \quad (15)$$

There are two special cases. When  $\rho \rightarrow +\infty$ , the space-time tends to the Minkowski space-time, and  $\sigma(\lambda) \rightarrow \lambda$ . When  $\rho = 1$ , the space-time becomes the AdS with curvature  $-1$ , and  $\sigma = \frac{\lambda - 2x_1}{x_2}$

### 3. Separation of Variables

Now we introduce a new Lax system

$$\partial_j \Psi = \sum_{\alpha=1}^r \frac{\sigma_{\alpha}^{j-1} B_{\alpha}}{\lambda - \lambda_{\alpha}} \Psi \quad (j = 1, 2, 3). \quad (16)$$

Here  $\lambda_1, \dots, \lambda_r$  are distinct complex numbers,  $\sigma_{\alpha} = \sigma(\lambda_{\alpha})$ ,  $B_1, \dots, B_r$  are  $N \times N$  matrix functions to be determined.

The integrability conditions of (16) are

$$\begin{aligned}& \partial_l(\sigma_{\alpha}^{k-1} B_{\alpha}) - \partial_k(\sigma_{\alpha}^{l-1} B_{\alpha}) \\ & + \sum_{\beta \neq \alpha} \frac{\sigma_{\alpha}^{k-1} \sigma_{\beta}^{l-1} - \sigma_{\alpha}^{l-1} \sigma_{\beta}^{k-1}}{\lambda_{\alpha} - \lambda_{\beta}} [B_{\alpha}, B_{\beta}] = 0\end{aligned}\quad (17)$$

( $1 \leq k < l \leq 3$ ). In fact, (17) is equivalent to two equations within it with  $k = 1, l = 2$  and  $k = 2, l = 3$  respectively:

$$\begin{aligned}\partial_2 B_{\alpha} &= \partial_1(\sigma_{\alpha} B_{\alpha}) + \left[ B_{\alpha}, \sum_{\beta=1}^r \gamma^{-1} B_{\beta} \right], \\ \partial_3(\sigma_{\alpha} B_{\alpha}) &= \partial_2(\sigma_{\alpha}^2 B_{\alpha}) + \left[ \sigma_{\alpha} B_{\alpha}, \sum_{\beta=1}^r \gamma^{-1} \sigma_{\beta} B_{\beta} \right].\end{aligned}\quad (18)$$

$B_{\alpha}$ 's can be constructed from the solutions of the original Lax pair (13) and its conjugate, as in the nonlinear constraint method for other equations.<sup>2</sup>

Suppose that  $F_\alpha$  is an  $N \times p$  solution ( $1 \leq p \leq N$ ) of (13) with  $\lambda = \lambda_\alpha$ , that is,  $F_\alpha$  satisfies

$$\begin{aligned}\partial_2 F_\alpha &= \sigma_\alpha \partial_1 F_\alpha - P F_\alpha, \\ \sigma_\alpha \partial_3 F_\alpha &= \sigma_\alpha^2 \partial_2 F_\alpha - \sigma_\alpha Q F_\alpha.\end{aligned}\quad (19)$$

Suppose also that  $G_\alpha$  is an  $N \times p$  solution of

$$\begin{aligned}\partial_2 G_\alpha &= \partial_1(\sigma_\alpha G_\alpha) + P^T G_\alpha, \\ \sigma_\alpha \partial_3 G_\alpha &= \partial_2(\sigma_\alpha^2 G_\alpha) + \sigma_\alpha Q^T G_\alpha,\end{aligned}\quad (20)$$

which is conjugate to (19).

Let  $B_\alpha = \kappa_\alpha F_\alpha G_\alpha^T$  where  $\kappa_\alpha$  is a complex constant, then  $B_\alpha$  satisfies

$$\begin{aligned}\partial_2 B_\alpha &= \partial_1(\sigma_\alpha B_\alpha) + [B_\alpha, P], \\ \partial_3(\sigma_\alpha B_\alpha) &= \partial_2(\sigma_\alpha^2 B_\alpha) + [\sigma_\alpha B_\alpha, Q].\end{aligned}\quad (21)$$

Comparing (21) with (18), we know that when

$$P = \gamma^{-1} \sum_{\beta=1}^r B_\beta, \quad Q = \gamma^{-1} \sum_{\beta=1}^r \sigma_\beta B_\beta, \quad (22)$$

(18) is consistent with (21).

Neglecting the special expressions of  $B_\alpha$ 's, we have the following general theorem.

**Theorem 1.** Suppose  $\lambda_1, \dots, \lambda_r$  are distinct complex constants, and  $B_1, \dots, B_r$  are  $N \times N$  matrices satisfying the integrability conditions (18) of (16), then

$$\begin{aligned}A_1 &= 0, \quad A_2 = \frac{1}{2} \gamma^{-1} \sum_{\beta=1}^r B_\beta, \\ A_3 &= \gamma^{-1} \sum_{\beta=1}^r \sigma_\beta B_\beta, \quad \Phi = \frac{1}{2} \sum_{\beta=1}^r B_\beta\end{aligned}\quad (23)$$

give a solution of the Bogomolny equation (8).

**Proof:** Since the gauge has already been fixed so that  $A_1 = 0$ ,  $A_2 = \gamma^{-1} \Phi$ , we have

$$D_1 \Phi = \gamma \partial_1 A_2 = \gamma F_{12}, \quad (24)$$

which is the first equation of (8). Now we verify the other two equations of (8). Summing up (18) on  $\alpha$ , we have

$$2D_2 \Phi = 2\partial_2 \Phi = \sum_{\alpha=1}^r \partial_2 B_\alpha = \gamma \partial_1 A_3 = \gamma F_{13}, \quad (25)$$

$$\begin{aligned}
 2D_3\Phi &= 2\partial_3\Phi + 2[A_3, \Phi] = \partial_3\left(\sum_{\alpha=1}^r B_\alpha\right) - \left[\sum_{\alpha=1}^r B_\alpha, \sum_{\beta=1}^r \gamma^{-1}\sigma_\beta B_\beta\right] \\
 &= \partial_2\left(\sum_{\alpha=1}^r \sigma_\alpha B_\alpha\right) + \sum_{\alpha=1}^r (\partial_2\sigma_\alpha)B_\alpha = \partial_2(\gamma A_3) - \rho^{-1}A_3 \\
 &= \gamma\partial_2A_3 = \gamma F_{23} + D_3\Phi.
 \end{aligned}
 \tag{26}$$

So  $D_3\Phi = \gamma F_{23}$ . The theorem is proved.

The solutions of (18) can be obtained by standard ways in 1+1 dimensions. In the next section, we will use the Darboux transformation method to solve them.

**Remark 1.** (16) can be generalized to  $j = 1, \dots, n$  with any positive integer  $n$ . For  $n = 2$ ,  $r = 2$ ,  $\lambda_1 = 1$ ,  $\lambda_2 = -1$ , and  $\rho \rightarrow +\infty$ , the Lax system is

$$\begin{aligned}
 \partial_1\Psi &= \left(\frac{B_1}{\lambda-1} + \frac{B_2}{\lambda+1}\right)\Psi, \\
 \partial_2\Psi &= \left(\frac{B_1}{\lambda-1} - \frac{B_2}{\lambda+1}\right)\Psi.
 \end{aligned}
 \tag{27}$$

This is the well-known Lax pair of the 1+1 dimensional principal chiral field (harmonic map from  $\mathbf{R}^{1,1}$  to the Lie group  $G$ ). A lot of explicit solutions of it have been known already.<sup>6,12,13,16</sup>

### 4. Darboux Transformation

We use the general procedure of constructing Darboux transformation for (16).<sup>4</sup>

Let  $\mu_1, \dots, \mu_N$  be non-zero complex numbers such that  $\mu_1, \dots, \mu_N$ ,  $\lambda_1, \dots, \lambda_r$  are distinct,  $h_j$  be a solution of (16) with  $\lambda = \mu_j$  ( $j = 1, \dots, N$ ). Let

$$\begin{aligned}
 \Lambda &= \text{diag}(\mu_1, \dots, \mu_N), \quad H = (h_1, \dots, h_N), \\
 S &= H\Lambda H^{-1}.
 \end{aligned}
 \tag{28}$$

Then, for any solution  $\Psi$  of (16),  $\tilde{\Psi} = (\lambda I - S)\Psi$  satisfies

$$\partial_j \tilde{\Psi} = \sum_{\alpha=1}^r \frac{\sigma_\alpha^{j-1} \tilde{B}_\alpha}{\lambda - \lambda_\alpha} \tilde{\Psi}
 \tag{29}$$

where

$$\tilde{B}_\alpha = (\lambda_\alpha I - S)B_\alpha(\lambda_\alpha I - S)^{-1}
 \tag{30}$$

and  $S$  satisfies

$$\partial_j S = \sum_{\alpha=1}^r \sigma_{\alpha}^{j-1} (B_{\alpha} - \tilde{B}_{\alpha}). \quad (31)$$

(31) can also be written as

$$\sum_{\alpha=1}^r \sigma_{\alpha}^{j-1} \tilde{B}_{\alpha} = \sum_{\alpha=1}^r \sigma_{\alpha}^{j-1} B_{\alpha} - \partial_j S. \quad (32)$$

From (23), this gives a Darboux transformation as stated in the following theorem.

**Theorem 2.** Suppose  $B_{\alpha}$ 's are solutions of (18),  $S$  is constructed as above,  $(A_1, A_2, A_3, \Phi)$  are given by (23), then

$$\begin{aligned} \tilde{A}_1 &= 0, & \tilde{A}_2 &= A_2 - \frac{\rho}{2(x_2 + \rho - 1)} \partial_1 S, \\ \tilde{A}_3 &= A_3 - \frac{\rho}{x_2 + \rho - 1} \partial_2 S, & \tilde{\Phi} &= \Phi - \frac{1}{2} \partial_1 S \end{aligned} \quad (33)$$

also give a solution of the Bogomolny equation (17).

This is the Darboux transformation of degree one. An infinite series of solutions can be obtained by successive actions of Darboux transformations of degree one.

**Example.** To use the Darboux transformation, we need a seed solution of (18). Suppose the seed solution is  $B_{\alpha} = u_{\alpha}(x_1, x_2, x_3)C_{\alpha}$  ( $\alpha = 1, \dots, r$ ) where  $u_{\alpha}$ 's are functions,  $C_{\alpha}$ 's are constant  $N \times N$  matrices such that  $[C_{\alpha}, C_{\beta}] = 0$  for  $\alpha, \beta = 1, \dots, r$ . By (18),  $u_{\alpha}$ 's satisfy

$$\begin{aligned} \partial_2 u_{\alpha} &= \partial_1 (\sigma_{\alpha} u_{\alpha}), \\ \partial_3 (\sigma_{\alpha} u_{\alpha}) &= \partial_2 (\sigma_{\alpha}^2 u_{\alpha}). \end{aligned} \quad (34)$$

The general solution is

$$u_{\alpha} = \left( \frac{x_2 + \rho - 1}{\rho \lambda_{\alpha} - 2x_1} \right)^2 f_{\alpha}(\theta_{\alpha}(x_1, x_2, x_3)) \quad (35)$$

where

$$\theta_{\alpha}(x_1, x_2, x_3) = \frac{(x_2 + \rho - 1)^2}{2(\rho \lambda_{\alpha} - 2x_1)} + x_3 - \frac{\rho - 2}{2\lambda_{\alpha}}, \quad (36)$$

$f_{\alpha}$ 's are arbitrary meromorphic functions.

**Remark 2.** In (36), the constant  $-\frac{\rho - 2}{2\lambda_{\alpha}}$  is added so that  $\theta_{\alpha}$  converges as  $\rho \rightarrow +\infty$ . Hence, in the Minkowski space-time,  $u_{\alpha} = \lambda_{\alpha}^{-2} f_{\alpha}(x_3 + \lambda_{\alpha}^{-1} x_2 + \lambda_{\alpha}^{-2} x_1)$ .

(35) gives the seed solution of (18) for the Darboux transformation, and the corresponding

$$\begin{aligned} A_1 &= 0, \quad A_2 = \frac{\rho}{2(x_2 + \rho - 1)} \sum_{\alpha=1}^r u_\alpha C_\alpha, \\ A_3 &= \sum_{\alpha=1}^r \frac{\rho(\rho\lambda_\alpha - 2x_1)}{(x_2 + \rho - 1)^2} u_\alpha C_\alpha, \quad \Phi = \frac{1}{2} \sum_{\alpha=1}^r u_\alpha C_\alpha. \end{aligned} \quad (37)$$

The fundamental solution of (16) is

$$\Psi = \exp \left( \sum_{\alpha=1}^r \frac{F_\alpha(\theta_\alpha) C_\alpha}{\lambda - \lambda_\alpha} \right) \Psi_0 \quad (38)$$

where  $\Psi_0$  is an  $N \times N$  constant matrix,  $F'_\alpha = f_\alpha$ .

With the seed solution (37), we can get new explicit solutions according to Theorem 2.

Especially, take  $G = SU(2)$ ,  $C_\alpha = \text{diag}(ic_\alpha, -ic_\alpha)$  where  $c_\alpha$ 's are constant real numbers. Then, as usual construction for  $SU(2)$  solutions, we should take  $\mu_2 = \bar{\mu}_1$  and  $h_2^* h_1 = 0$ . Hence

$$H = \begin{pmatrix} \alpha_1 g & -\bar{\alpha}_2 \bar{g}^{-1} \\ \alpha_2 g^{-1} & \bar{\alpha}_1 \bar{g} \end{pmatrix} \quad (39)$$

where

$$g = \exp \left( i \sum_{\alpha=1}^r \frac{F_\alpha(\theta_\alpha) c_\alpha}{\mu_1 - \lambda_\alpha} \right), \quad (40)$$

$\alpha_1$  and  $\alpha_2$  are complex constants.

$$\begin{aligned} S &= H \Lambda H^{-1} \\ &= \frac{\mu_1 + \bar{\mu}_1}{2} I + \frac{\mu_1 - \bar{\mu}_1}{2} \frac{1}{|\alpha_1|^2 |g|^2 + |\alpha_2|^2 |g|^{-2}} \\ &\quad \cdot \begin{pmatrix} |\alpha_1|^2 |g|^2 - |\alpha_2|^2 |g|^{-2} & 2\alpha_1 \bar{\alpha}_2 g \bar{g}^{-1} \\ 2\bar{\alpha}_1 \alpha_2 \bar{g} g^{-1} & -|\alpha_1|^2 |g|^2 + |\alpha_2|^2 |g|^{-2} \end{pmatrix}, \end{aligned} \quad (41)$$

and

$$\begin{aligned} \tilde{A}_1 &= 0, \\ \tilde{A}_2 &= \frac{\rho}{2(x_2 + \rho - 1)} \sum_{\alpha=1}^r u_\alpha C_\alpha - \frac{\rho}{2(x_2 + \rho - 1)} \partial_1 S, \\ \tilde{A}_3 &= \sum_{\alpha=1}^r \frac{\rho(\rho\lambda_\beta - 2x_1)}{(x_2 + \rho - 1)^2} u_\alpha C_\alpha - \frac{\rho}{x_2 + \rho - 1} \partial_2 S, \\ \tilde{\Phi} &= \frac{1}{2} \sum_{\alpha=1}^r u_\alpha C_\alpha - \frac{1}{2} \partial_1 S. \end{aligned} \quad (42)$$

## References

1. C.W. Cao, Y.T. Wu and X.G. Geng, Relation between the Kadomtsev-Petviashvili equation and the confocal involutive system, *J. Math. Phys.* **40** (1999) 3948-3970.
2. Y. Cheng and Y.S. Li, The constraint of the KP equation and its special solutions, *Phys. Lett.* **A157** (1991) 22-26.
3. B. Dai and C.L. Terng, Backlund transformations, Ward solitons, and unitons, *J. Diff. Geom.* **75** (2007) 57-108.
4. C.H. Gu, On the Darboux form of Bäcklund transformations, in: X.C. Song (Eds.), *Integrable System*, World Scientific (1989) 162-168.
5. C.H. Gu, Darboux transformation and solitons of Yang-Mills-Higgs equations in  $R^{2,1}$ , *Science in China* **A45** (2002) 706-715.
6. C.H. Gu and H.S. Hu, Construction of unitons via purely algebraic algorithm, *Chin. Ann. Math.* **20B** (1999) 1-6.
7. C.H. Gu and Z.X. Zhou, On Darboux transformations for soliton equations in high dimensional space-time, *Lett. Math. Phys.* **32** (1994) 1-10.
8. N.J. Hitchin, G.B. Segal and R.S. Ward: *Integrable systems, Twistors, loop groups and Riemann surfaces*, (Clarendon Press, Oxford), 1999.
9. T. Ioannidou, Bogomolny Yang-Mills-Higgs solutions in (2+1) anti-de Sitter space, *Nonlinearity* **15** (2002) 1489-1497.
10. T. Ioannidou, Nontrivial soliton scattering in planar integrable systems, *Intl. J. Mod. Phys.* **A18** (2003) 4975-4998.
11. B. Konopelchenko, J. Sidorenko and W. Strampp, (1+1)-dimensional integrable systems as symmetry constraints of (2+1) dimensional systems, *Phys. Lett.* **A157** (1991) 17-21.
12. A.T. Ogielski, K. Prasad, A. Sinha and L.L. Chau, Bäcklund transformations and local conservation laws for principal chiral fields, *Phys. Lett.* **B91** (1980) 387-391.
13. K. Uhlenbeck, Harmonic maps into Lie groups: classical solutions of the chiral model, *J. Diff. Geom.* **38** (1980) 1-50.
14. R.S. Ward, Soliton solutions in an integrable chiral model in 2+1 dimensions, *J. Math. Phys.* **29** (1988) 386-389.
15. R.S. Ward, Two integrable systems related to hyperbolic monopoles, *Asian J. Math.* **3** (1999) 325-332.
16. V.E. Zakharov and A.V. Mikhailov, Relativistically invariant two-dimensional models of field theory which are integrable by means of the inverse problem scattering method, *Sov. Phys. JETP* **47** (1978) 1017-1027.
17. Z.X. Zhou, Solutions of the Yang-Mills-Higgs equations in 2+1-dimensional anti-de Sitter space-time, *J. Math. Phys.* **42** (2001) 1085-1099.
18. Z.X. Zhou, Relation between the solitons of Yang-Mills-Higgs fields in 2+1 dimensional Minkowski space-time and anti-de Sitter space-time, *J. Math. Phys.* **42** (2001) 4938-4946.
19. Z.X. Zhou, Localized solitons of hyperbolic  $su(N)$  AKNS system, *Inverse Problems* **14** (1998) 1371-1383.

20. Z.X. Zhou, W.X. Ma and R.G. Zhou, Finite-dimensional integrable systems associated with Davey-Stewartson I equation, *Nonlinearity* **14** (2001) 701-717.