

KCs where the old and new claw methods differ most (page 1 of 5)
Each KC: left = old claws (2 µm straight-line single linkage on PN→KC PSDs); right = new claws (Yang's rule of 2026-11-04: all calyx synapse sites clustered by single linkage at 8 µm along the KC's skeleton; each PN with ≥2 PSD(s) in a cluster gets its own claw; other sites join the claw of the nearest PN→KC PSD if it is ≤8 µm away along the skeleton).
Big symbols = PN→KC PSDs; color + shape = glomerulus of the presynaptic PN (the same in both panels; hollow = PSD not on a claw).
Small dots = the KC's other calyx synapse sites, colored by the glomerulus of the claw they join (grey = not on a claw). Thin outline = one claw.
Label = glomerulus of the claw's main PN (claws with ≥3 PN PSDs); * = claw sharing its 8 µm cluster with another PN's claw. Grey lines = KC skeleton near its calyx sites.
View = the plane in which the KC's calyx sites spread most (same for both panels).


hemibrain R · 600011602 · KCab-m · 78 PN PSDs · Prat 5 (corrected 5)
old: 2 µm, PN PSDs: 15 claws




new: 8 µm along skeleton, split by PN: 5 claws




hemibrain R · 736808904 · KCab-s · 122 PN PSDs · Prat 7 (corrected 8)
old: 2 µm, PN PSDs: 15 claws




new: 8 µm along skeleton, split by PN: 7 claws




hemibrain R · 5813021313 · KCab-m · 74 PN PSDs · Prat 6 (corrected 7)
old: 2 µm, PN PSDs: 12 claws




new: 8 µm along skeleton, split by PN: 5 claws




hemibrain R · 569943419 · KCab-m · 90 PN PSDs · Prat 5 (corrected 5)
old: 2 µm, PN PSDs: 11 claws




new: 8 µm along skeleton, split by PN: 4 claws




hemibrain R · 363029976 · KCab-m · 88 PN PSDs · Prat 6 (corrected 7)
old: 2 µm, PN PSDs: 14 claws




new: 8 µm along skeleton, split by PN: 7 claws




hemibrain R · 5812981673 · KCab-s · 100 PN PSDs · Prat 7 (corrected 7)
old: 2 µm, PN PSDs: 14 claws




new: 8 µm along skeleton, split by PN: 7 claws




hemibrain R · 939742609 · KCab-c · 59 PN PSDs · Prat 4 (corrected 6)
old: 2 µm, PN PSDs: 13 claws




new: 8 µm along skeleton, split by PN: 6 claws



hemibrain R · 5812981071 · KCa'b'-m · 82 PN PSDs · Prat 4 (corrected 6)
old: 2 µm, PN PSDs: 18 claws




new: 8 µm along skeleton, split by PN: 6 claws




KCs where the old and new claw methods differ most (page 2 of 5)
Each KC: left = old claws (2 µm straight-line single linkage on PN→KC PSDs); right = new claws (Yang's rule of 2026-11-04: all calyx synapse sites clustered by single linkage at 8 µm along the KC's skeleton; each PN with ≥2 PSD(s) in a cluster gets its own claw; other sites join the claw of the nearest PN→KC PSD if it is ≤8 µm away along the skeleton).
Big symbols = PN→KC PSDs; color + shape = glomerulus of the presynaptic PN (the same in both panels; hollow = PSD not on a claw).
Small dots = the KC's other calyx synapse sites, colored by the glomerulus of the claw they join (grey = not on a claw). Thin outline = one claw.
Label = glomerulus of the claw's main PN (claws with ≥3 PN PSDs); * = claw sharing its 8 µm cluster with another PN's claw. Grey lines = KC skeleton near its calyx sites.
View = the plane in which the KC's calyx sites spread most (same for both panels).


hemibrain R · 673029543 · KCa'b'-m · 63 PN PSDs · Prat 7 (corrected 6)
old: 2 µm, PN PSDs: 19 claws




new: 8 µm along skeleton, split by PN: 8 claws




hemibrain R · 5812982692 · KCa'b'-m · 64 PN PSDs · Prat 4 (corrected 4)
old: 2 µm, PN PSDs: 13 claws




new: 8 µm along skeleton, split by PN: 4 claws




hemibrain R · 5812982706 · KCa'b'-m · 82 PN PSDs · Prat 5 (corrected 6)
old: 2 µm, PN PSDs: 17 claws




new: 8 µm along skeleton, split by PN: 8 claws




hemibrain R · 5812982329 · KCa'b'-m · 85 PN PSDs · Prat 5 (corrected 5)
old: 2 µm, PN PSDs: 14 claws




new: 8 µm along skeleton, split by PN: 6 claws




hemibrain R · 548552875 · KCa'b'-m · 54 PN PSDs · Prat 4 (corrected 4)
old: 2 µm, PN PSDs: 13 claws




new: 8 µm along skeleton, split by PN: 5 claws




hemibrain R · 661430106 · KCg-m · 168 PN PSDs · Prat 10 (corrected 10)
old: 2 µm, PN PSDs: 20 claws




new: 8 µm along skeleton, split by PN: 10 claws




hemibrain R · 5813084692 · KCg-m · 192 PN PSDs · Prat 10 (corrected 11)
old: 2 µm, PN PSDs: 18 claws




new: 8 µm along skeleton, split by PN: 9 claws



hemibrain R · 785232649 · KCg-m · 127 PN PSDs · Prat 6 (corrected 8)
old: 2 µm, PN PSDs: 16 claws




new: 8 µm along skeleton, split by PN: 7 claws




KCs where the old and new claw methods differ most (page 3 of 5)
Each KC: left = old claws (2 µm straight-line single linkage on PN→KC PSDs); right = new claws (Yang's rule of 2026-11-04: all calyx synapse sites clustered by single linkage at 8 µm along the KC's skeleton; each PN with ≥2 PSD(s) in a cluster gets its own claw; other sites join the claw of the nearest PN→KC PSD if it is ≤8 µm away along the skeleton).
Big symbols = PN→KC PSDs; color + shape = glomerulus of the presynaptic PN (the same in both panels; hollow = PSD not on a claw).
Small dots = the KC's other calyx synapse sites, colored by the glomerulus of the claw they join (grey = not on a claw). Thin outline = one claw.
Label = glomerulus of the claw's main PN (claws with ≥3 PN PSDs); * = claw sharing its 8 µm cluster with another PN's claw. Grey lines = KC skeleton near its calyx sites.
View = the plane in which the KC's calyx sites spread most (same for both panels).


hemibrain R · 847302850 · KCg-m · 154 PN PSDs · Prat 8 (corrected 11)
old: 2 µm, PN PSDs: 19 claws




new: 8 µm along skeleton, split by PN: 10 claws




hemibrain R · 632082635 · KCg-m · 199 PN PSDs · Prat 12 (corrected 12)
old: 2 µm, PN PSDs: 21 claws




new: 8 µm along skeleton, split by PN: 12 claws




hemibrain R · 1173745411 · KCg-m · 163 PN PSDs · Prat 9 (corrected 9)
old: 2 µm, PN PSDs: 18 claws




new: 8 µm along skeleton, split by PN: 9 claws




hemibrain R · 723495119 · KCg-m · 144 PN PSDs · Prat 10 (corrected 12)
old: 2 µm, PN PSDs: 19 claws




new: 8 µm along skeleton, split by PN: 10 claws




male CNS L · 57902 · KCab-s · 94 PN PSDs
old: 2 µm, PN PSDs: 15 claws




new: 8 µm along skeleton, split by PN: 6 claws




male CNS L · 168916 · KCab-m · 94 PN PSDs
old: 2 µm, PN PSDs: 12 claws




new: 8 µm along skeleton, split by PN: 4 claws




male CNS L · 143895 · KCab-m · 123 PN PSDs
old: 2 µm, PN PSDs: 13 claws




new: 8 µm along skeleton, split by PN: 5 claws



male CNS L · 82480 · KCa'b'-m · 88 PN PSDs
old: 2 µm, PN PSDs: 13 claws




new: 8 µm along skeleton, split by PN: 3 claws




KCs where the old and new claw methods differ most (page 4 of 5)
Each KC: left = old claws (2 µm straight-line single linkage on PN→KC PSDs); right = new claws (Yang's rule of 2026-11-04: all calyx synapse sites clustered by single linkage at 8 µm along the KC's skeleton; each PN with ≥2 PSD(s) in a cluster gets its own claw; other sites join the claw of the nearest PN→KC PSD if it is ≤8 µm away along the skeleton).
Big symbols = PN→KC PSDs; color + shape = glomerulus of the presynaptic PN (the same in both panels; hollow = PSD not on a claw).
Small dots = the KC's other calyx synapse sites, colored by the glomerulus of the claw they join (grey = not on a claw). Thin outline = one claw.
Label = glomerulus of the claw's main PN (claws with ≥3 PN PSDs); * = claw sharing its 8 µm cluster with another PN's claw. Grey lines = KC skeleton near its calyx sites.
View = the plane in which the KC's calyx sites spread most (same for both panels).


male CNS L · 74531 · KCa'b'-m · 116 PN PSDs
old: 2 µm, PN PSDs: 19 claws




new: 8 µm along skeleton, split by PN: 9 claws




male CNS L · 84602 · KCa'b'-m · 60 PN PSDs
old: 2 µm, PN PSDs: 16 claws




new: 8 µm along skeleton, split by PN: 7 claws




male CNS L · 69417 · KCg-m · 140 PN PSDs
old: 2 µm, PN PSDs: 15 claws




new: 8 µm along skeleton, split by PN: 5 claws




male CNS L · 75597 · KCg-m · 145 PN PSDs
old: 2 µm, PN PSDs: 16 claws




new: 8 µm along skeleton, split by PN: 7 claws




male CNS L · 135480 · KCg-m · 72 PN PSDs
old: 2 µm, PN PSDs: 13 claws




new: 8 µm along skeleton, split by PN: 5 claws




male CNS L · 79783 · KCg-m · 102 PN PSDs
old: 2 µm, PN PSDs: 14 claws




new: 8 µm along skeleton, split by PN: 6 claws




male CNS R · 106981 · KCab-s · 67 PN PSDs
old: 2 µm, PN PSDs: 13 claws




new: 8 µm along skeleton, split by PN: 4 claws



male CNS R · 80172 · KCab-s · 105 PN PSDs
old: 2 µm, PN PSDs: 17 claws




new: 8 µm along skeleton, split by PN: 7 claws




KCs where the old and new claw methods differ most (page 5 of 5)
Each KC: left = old claws (2 µm straight-line single linkage on PN→KC PSDs); right = new claws (Yang's rule of 2026-11-04: all calyx synapse sites clustered by single linkage at 8 µm along the KC's skeleton; each PN with ≥2 PSD(s) in a cluster gets its own claw; other sites join the claw of the nearest PN→KC PSD if it is ≤8 µm away along the skeleton).
Big symbols = PN→KC PSDs; color + shape = glomerulus of the presynaptic PN (the same in both panels; hollow = PSD not on a claw).
Small dots = the KC's other calyx synapse sites, colored by the glomerulus of the claw they join (grey = not on a claw). Thin outline = one claw.
Label = glomerulus of the claw's main PN (claws with ≥3 PN PSDs); * = claw sharing its 8 µm cluster with another PN's claw. Grey lines = KC skeleton near its calyx sites.
View = the plane in which the KC's calyx sites spread most (same for both panels).


male CNS R · 74869 · KCab-s · 110 PN PSDs
old: 2 µm, PN PSDs: 15 claws




new: 8 µm along skeleton, split by PN: 7 claws




male CNS R · 71908 · KCa'b'-m · 74 PN PSDs
old: 2 µm, PN PSDs: 19 claws




new: 8 µm along skeleton, split by PN: 6 claws




male CNS R · 64782 · KCa'b'-ap2 · 62 PN PSDs
old: 2 µm, PN PSDs: 16 claws




new: 8 µm along skeleton, split by PN: 6 claws




male CNS R · 82744 · KCa'b'-m · 90 PN PSDs
old: 2 µm, PN PSDs: 15 claws




new: 8 µm along skeleton, split by PN: 5 claws




male CNS R · 149792 · KCg-m · 129 PN PSDs
old: 2 µm, PN PSDs: 19 claws




new: 8 µm along skeleton, split by PN: 8 claws




male CNS R · 75029 · KCg-m · 215 PN PSDs
old: 2 µm, PN PSDs: 36 claws




new: 8 µm along skeleton, split by PN: 26 claws




male CNS R · 58014 · KCg-m · 89 PN PSDs
old: 2 µm, PN PSDs: 14 claws



new: 8 µm along skeleton, split by PN: 5 claws



male CNS R · 68226 · KCg-m · 124 PN PSDs
old: 2 µm, PN PSDs: 15 claws



new: 8 µm along skeleton, split by PN: 6 claws

