

The Effect of Strain on Charge Density Wave Order in the Holstein Model

Benjamin Cohen-Stead

May 13, 2019

Outline for Talk

- 1 Background
- 2 The Holstein Model
- 3 Charge-Density Wave Order
- 4 Describing My Project
- 5 Review DQMC (Determinant Quantum Monte Carlo)
- 6 Results
- 7 Summary

Background

Interactions in condensed matter systems can give rise to strongly correlated systems, resulting in novel electronic phases.

Background

Interactions in condensed matter systems can give rise to strongly correlated systems, resulting in novel electronic phases.

Types of Interactions:

- Electron-Electron Interactions
- Electron-Ion Interactions

⇒ Electron-Phonon Inter.

Background

Interactions in condensed matter systems can give rise to strongly correlated systems, resulting in novel electronic phases.

Types of Interactions:

- Electron-Electron Interactions
- Electron-Ion Interactions
- ⇒ Electron-Phonon Inter.

Emergent Phases in Solids:

- (Anti)Ferromagnetism
- Superconductivity
- Charge-Density Wave Order (CDW)

Background

Interactions in condensed matter systems can give rise to strongly correlated systems, resulting in novel electronic phases.

Types of Interactions:

- Electron-Electron Interactions
- Electron-Ion Interactions
- ⇒ Electron-Phonon Inter.

Emergent Phases in Solids:

- (Anti)Ferromagnetism
- Superconductivity
- Charge-Density Wave Order (CDW)

Model Hamiltonians:

- Simple ⇒ small number of parameters
- Describes most important interactions
- Reproduces underlying physics of emergent phases
- Does not incorporate details of real materials

The Holstein Model

The Holstein Model: A model Hamiltonian for describing electron-phonon interactions on a lattice.

$$\hat{H} = \hat{K} + \hat{U} + \hat{V}$$

The Holstein Model

The Holstein Model: A model Hamiltonian for describing electron-phonon interactions on a lattice.

$$\hat{H} = \hat{K} + \hat{U} + \hat{V}$$

- **Electron Kinetic Energy:** $\hat{T} = -t \sum_{\langle i,j \rangle \sigma} \left(\hat{c}_{i,\sigma}^\dagger \hat{c}_{j,\sigma} + h.c. \right) - \mu \sum_{i,\sigma} \hat{n}_{i,\sigma}$

The Holstein Model

The Holstein Model: A model Hamiltonian for describing electron-phonon interactions on a lattice.

$$\hat{H} = \hat{K} + \hat{U} + \hat{V}$$

- **Electron Kinetic Energy:** $\hat{T} = -t \sum_{\langle i,j \rangle \sigma} \left(\hat{c}_{i,\sigma}^\dagger \hat{c}_{j,\sigma} + h.c. \right) - \mu \sum_{i,\sigma} \hat{n}_{i,\sigma}$
- **Bare Phonon Term:** $\hat{U} = \frac{\omega_0^2}{2} \sum_i \hat{X}_i^2 + \frac{1}{2} \sum_i \hat{P}_i^2$

The Holstein Model

The Holstein Model: A model Hamiltonian for describing electron-phonon interactions on a lattice.

$$\hat{H} = \hat{K} + \hat{U} + \hat{V}$$

- **Electron Kinetic Energy:** $\hat{T} = -t \sum_{\langle i,j \rangle \sigma} \left(\hat{c}_{i,\sigma}^\dagger \hat{c}_{j,\sigma} + h.c. \right) - \mu \sum_{i,\sigma} \hat{n}_{i,\sigma}$
- **Bare Phonon Term:** $\hat{U} = \frac{\omega_0^2}{2} \sum_i \hat{X}_i^2 + \frac{1}{2} \sum_i \hat{P}_i^2$
- **Electron-Phonon Interaction:** $\hat{V} = \lambda \sum_{i,\sigma} \hat{X}_i \hat{n}_{i,\sigma}$

The Holstein Model

The Holstein Model: A model Hamiltonian for describing electron-phonon interactions on a lattice.

$$\hat{H} = \hat{K} + \hat{U} + \hat{V}$$

- **Electron Kinetic Energy:** $\hat{T} = -t \sum_{\langle i,j \rangle \sigma} \left(\hat{c}_{i,\sigma}^\dagger \hat{c}_{j,\sigma} + h.c. \right) - \mu \sum_{i,\sigma} \hat{n}_{i,\sigma}$
- **Bare Phonon Term:** $\hat{U} = \frac{\omega_0^2}{2} \sum_i \hat{X}_i^2 + \frac{1}{2} \sum_i \hat{P}_i^2$
- **Electron-Phonon Interaction:** $\hat{V} = \lambda \sum_{i,\sigma} \hat{X}_i \hat{n}_{i,\sigma}$
- **Half-Filling:** $\mu = -\frac{\lambda^2}{\omega_0^2} \Rightarrow \langle n \rangle = \langle n_\uparrow + n_\downarrow \rangle = 1$

The Holstein Model

The Holstein Model: A model Hamiltonian for describing electron-phonon interactions on a lattice.

$$\hat{H} = \hat{K} + \hat{U} + \hat{V}$$

- **Electron Kinetic Energy:** $\hat{T} = -t \sum_{\langle i,j \rangle \sigma} \left(\hat{c}_{i,\sigma}^\dagger \hat{c}_{j,\sigma} + h.c. \right) - \mu \sum_{i,\sigma} \hat{n}_{i,\sigma}$
- **Bare Phonon Term:** $\hat{U} = \frac{\omega_0^2}{2} \sum_i \hat{X}_i^2 + \frac{1}{2} \sum_i \hat{P}_i^2$
- **Electron-Phonon Interaction:** $\hat{V} = \lambda \sum_{i,\sigma} \hat{X}_i \hat{n}_{i,\sigma}$
- **Half-Filling:** $\mu = -\frac{\lambda^2}{\omega_0^2} \Rightarrow \langle n \rangle = \langle n_\uparrow + n_\downarrow \rangle = 1$
- **Dimensionless Coupling Constant:** $\lambda_D = \frac{\lambda^2}{\omega_0^2 W}$ where $W \sim t$ is bandwidth


The Holstein Electron-Phonon Interaction Term

Why is $\hat{V} = \lambda \sum_{i,\sigma} \hat{X}_i \hat{n}_{i,\sigma}$ a reasonable choice for an Electron-Phonon Interaction?

The Holstein Electron-Phonon Interaction Term

Why is $\hat{V} = \lambda \sum_{i,\sigma} \hat{X}_i \hat{n}_{i,\sigma}$ a reasonable choice for an Electron-Phonon Interaction?

$$\hat{V} = \frac{g}{\sqrt{N}} \sum_{k,q,\sigma} \left[\hat{b}_q^\dagger \hat{c}_{k-q,\sigma}^\dagger \hat{c}_{k,\sigma} + \hat{b}_q \hat{c}_{k+q,\sigma}^\dagger \hat{c}_{k,\sigma} \right], \quad g = \frac{\lambda}{\sqrt{2\omega_0}}$$



The Single-Site $t = 0$ Holstein Model at Half-filling

The Single-Site $t = 0$ Holstein Model at Half-filling

$$\hat{H} = \frac{1}{2}\hat{P}^2 + \frac{\omega_0^2}{2}\hat{X}^2 + \lambda\hat{X}\hat{n} + \frac{\lambda^2}{\omega_0^2}\hat{n}$$

The Single-Site $t = 0$ Holstein Model at Half-filling

$$\hat{H} = \frac{1}{2}\hat{P}^2 + \frac{\omega_0^2}{2}\hat{X}^2 + \lambda\hat{X}\hat{n} + \frac{\lambda^2}{\omega_0^2}\hat{n}$$

$$\hat{H} = \left[\frac{1}{2}\hat{P}^2 + \frac{\omega_0^2}{2} \left(\hat{X} + \frac{\lambda}{\omega_0^2}(\hat{n}_\uparrow + \hat{n}_\downarrow) \right)^2 \right]$$

$$+ \left[-\frac{\lambda^2}{2\omega_0^2} (2\hat{n}_\uparrow\hat{n}_\downarrow - \hat{n}_\uparrow - \hat{n}_\downarrow) \right]$$

The Single-Site $t = 0$ Holstein Model at Half-filling

$$\hat{H} = \frac{1}{2}\hat{P}^2 + \frac{\omega_0^2}{2}\hat{X}^2 + \lambda\hat{X}\hat{n} + \frac{\lambda^2}{\omega_0^2}\hat{n}$$

$$\hat{H} = \left[\frac{1}{2}\hat{P}^2 + \frac{\omega_0^2}{2} \left(\hat{X} + \frac{\lambda}{\omega_0^2}(\hat{n}_\uparrow + \hat{n}_\downarrow) \right)^2 \right]$$

$$+ \left[-\frac{\lambda^2}{2\omega_0^2} (2\hat{n}_\uparrow\hat{n}_\downarrow - \hat{n}_\uparrow - \hat{n}_\downarrow) \right]$$

- $|0\rangle \implies 0$
- $|\cdot\downarrow\rangle, |\uparrow\cdot\rangle \implies +\frac{\lambda^2}{2\omega_0^2}$
- $|\uparrow\downarrow\rangle \implies 0$

The Single-Site $t = 0$ Holstein Model at Half-filling

$$\hat{H} = \frac{1}{2}\hat{P}^2 + \frac{\omega_0^2}{2}\hat{X}^2 + \lambda\hat{X}\hat{n} + \frac{\lambda^2}{\omega_0^2}\hat{n} \implies U_{eff} = -\frac{\lambda^2}{\omega_0^2}$$

$$\hat{H} = \left[\frac{1}{2}\hat{P}^2 + \frac{\omega_0^2}{2} \left(\hat{X} + \frac{\lambda}{\omega_0^2}(\hat{n}_\uparrow + \hat{n}_\downarrow) \right)^2 \right]$$

$$+ \left[-\frac{\lambda^2}{2\omega_0^2} (2\hat{n}_\uparrow\hat{n}_\downarrow - \hat{n}_\uparrow - \hat{n}_\downarrow) \right]$$

- $|0\rangle \implies 0$
- $|\cdot\downarrow\rangle, |\uparrow\cdot\rangle \implies +\frac{\lambda^2}{2\omega_0^2}$
- $|\uparrow\downarrow\rangle \implies 0$


The Single-Site $t = 0$ Holstein Model at Half-filling

$$\hat{H} = \frac{1}{2}\hat{P}^2 + \frac{\omega_0^2}{2}\hat{X}^2 + \lambda\hat{X}\hat{n} + \frac{\lambda^2}{\omega_0^2}\hat{n}$$
$$\implies U_{\text{eff}} = -\frac{\lambda^2}{\omega_0^2}$$

$$\hat{H} = \left[\frac{1}{2}\hat{P}^2 + \frac{\omega_0^2}{2} \left(\hat{X} + \frac{\lambda}{\omega_0^2}(\hat{n}_\uparrow + \hat{n}_\downarrow) \right)^2 \right]$$

$$+ \left[-\frac{\lambda^2}{2\omega_0^2} (2\hat{n}_\uparrow\hat{n}_\downarrow - \hat{n}_\uparrow - \hat{n}_\downarrow) \right]$$

- $|0\rangle \implies 0$
- $|\cdot\downarrow\rangle, |\uparrow\cdot\rangle \implies +\frac{\lambda^2}{2\omega_0^2}$
- $|\uparrow\downarrow\rangle \implies 0$




Charge-Density Wave Order


Effective Electron-Electron Attraction Mediated by Phonons

⇒ Charge-Density Wave Order:

Periodic Spatial Modulation of the Electron Density



Charge-Density Wave Order




Peierls Instability

In a 1D chain, a uniform distribution of non-interacting electrons is unstable.

$$\chi_0(\mathbf{q}) = \sum_{\mathbf{k}} \frac{f(\epsilon_{\mathbf{k}}) - f(\epsilon_{\mathbf{k}+\mathbf{q}})}{\epsilon_{\mathbf{k}} - \epsilon_{\mathbf{k}+\mathbf{q}}}$$

Charge-Density Wave Order



Peierls Instability


In a 1D chain, a uniform distribution of non-interacting electrons is unstable.

$$\chi_0(\mathbf{q}) = \sum_{\mathbf{k}} \frac{f(\epsilon_{\mathbf{k}}) - f(\epsilon_{\mathbf{k}+\mathbf{q}})}{\epsilon_{\mathbf{k}} - \epsilon_{\mathbf{k}+\mathbf{q}}}$$

Fermi Surface Nesting allows Peierls explanation of CDW to be applied in $D > 1$ dimensions.

Charge-Density Wave Order

An example of a Peierls CDW in the quasi-1D material TTF-TCNQ
(tetrathiafulvalene-tetracyanoquinodimethane)



X. Zhu, J. Guo, J. Zhang, E. W. Plummer, (2017). Advances in Physics: X, 2(3), 622-640.

Charge-Density Wave Order

The Peierls Instability does not explain all CDW order in $D > 1$ dimensions.

- No FSN
- No Phonon Softening
- No Peak in χ_0
- No Metal-Insulator Transition

Charge-Density Wave Order



The Peierls's Instability does not explain all CDW order in $D > 1$ dimensions.

- No FSN
- No Phonon Softening
- No Peak in χ_0
- No Metal-Insulator Transition

Ex: 2H – NbSe₂

(Niobium diselenide)

- Quasi-2D System
- $g(\mathbf{q}) \Rightarrow$ CDW at $T_{\text{cdw}} \approx 33.5\text{K}$
- $q_{\text{cdw}} = 0.7\text{\AA}^{-1}$ ($a = 3.443\text{\AA}$)



The Project: A Strained Holstein Model

Effect of Strain on Charge Density Wave Order in the Holstein Model

B. Cohen-Stead,¹ N.C. Costa,^{2,3} E. Khatami,⁴ and R.T. Scalettar¹

¹*Department of Physics, University of California, Davis, CA 95616, USA**

²*International School for Advanced Studies (SISSA), Via Bonomea 265, 34136, Trieste, Italy*

³*Instituto de Física, Universidade Federal do Rio de Janeiro Cx.P. 68.528, 21941-972 Rio de Janeiro RJ, Brazil*

⁴*Department of Physics and Astronomy, San José State University, San José, CA 95192*

(Dated: May 7, 2019)

We investigate charge ordering in the Holstein model in the presence of anisotropic hopping, $t_x, t_y = 1 - \delta, 1 + \delta$, as a model of the effect of strain on charge density wave (CDW) materials. Using Quantum Monte Carlo simulations, we show that the CDW transition temperature is relatively insensitive to moderate anisotropy $\delta \lesssim 0.3$, but begins to decrease more rapidly at $\delta \gtrsim 0.4$. However, the density correlations, as well as the kinetic energies parallel and perpendicular to the compressional axis, change significantly for moderate δ . Accompanying mean-field theory calculations show a similar qualitative structure, with the transition temperature relatively constant at small δ and a more rapid decrease for larger strains. We also obtain the density of states $N(\omega)$, which provides clear signal of the charge ordering transition at large strain, where finite size scaling of the charge structure factor is extremely difficult because of the small value of the order parameter.

PACS numbers: 71.10.Fd, 71.30.+h, 71.45.Lr, 74.20.-z, 02.70.Uu

The Project: A Strained Holstein Model

The Strained Holstein Model:

- Square Lattice

The Project: A Strained Holstein Model

The Strained Holstein Model:

- Square Lattice
- $t_x = (1 - \delta)t$, $t_y = (1 + \delta)t$

The Project: A Strained Holstein Model

The Strained Holstein Model:

- Square Lattice
- $t_x = (1 - \delta)t, t_y = (1 + \delta)t$
- $t = 1 \implies W = 8t = 8$

The Project: A Strained Holstein Model



The Strained Holstein Model:

- Square Lattice
- $t_x = (1 - \delta)t, t_y = (1 + \delta)t$
- $t = 1 \implies W = 8t = 8$
- $\lambda = \sqrt{2}, \omega_0 = 1 \implies \lambda_D = 0.25$

The Project: A Strained Holstein Model

The Strained Holstein Model:

- Square Lattice
- $t_x = (1 - \delta)t, t_y = (1 + \delta)t$
- $t = 1 \implies W = 8t = 8$
- $\lambda = \sqrt{2}, \omega_0 = 1 \implies \lambda_D = 0.25$
- $\mathbf{q}_{\text{cdw}} = (\pi, \pi) \implies \text{Checkerboard}$



A Mean-Field Approximation

1. Adiabatic Approx. \Rightarrow Drop $\frac{1}{2} \sum_i \hat{P}_i^2$

A Mean-Field Approximation

1. Adiabatic Approx. \Rightarrow Drop $\frac{1}{2} \sum_i \hat{P}_i^2$

2. *Ansatz*: $\hat{X}_i \rightarrow x_0 + (-1)^i x_1$, where

$$x_0 = -\frac{\lambda}{\omega_0^2} \text{ and } x_1 = \text{order parameter}$$

A Mean-Field Approximation

1. Adiabatic Approx. \Rightarrow Drop $\frac{1}{2} \sum_i \hat{P}_i^2$

2. *Ansatz*: $\hat{X}_i \rightarrow x_0 + (-1)^i x_1$, where

$x_0 = -\frac{\lambda}{\omega_0^2}$ and x_1 = order parameter

3. Minimize the Free Energy:

$$F = \frac{N}{2} \omega_0^2 (x_0^2 + x_1^2) - \frac{2}{\beta} \sum_{\mathbf{k}} \ln(1 + e^{-\beta E_{\mathbf{k}}(x_1)})$$

A Mean-Field Approximation

1. Adiabatic Approx. \Rightarrow Drop $\frac{1}{2} \sum_i \hat{P}_i^2$

2. Ansatz: $\hat{X}_i \rightarrow x_0 + (-1)^i x_1$, where

$x_0 = -\frac{\lambda}{\omega_0^2}$ and $x_1 = \text{order parameter}$

3. Minimize the Free Energy:

$$F = \frac{N}{2} \omega_0^2 (x_0^2 + x_1^2) - \frac{2}{\beta} \sum_{\mathbf{k}} \ln(1 + e^{-\beta E_{\mathbf{k}}(x_1)})$$

4. $x_1 = 0$ corresponds to $T > T_{\text{cdw}}$

$x_1 > 0$ corresponds to $T < T_{\text{cdw}}$

A Mean-Field Approximation

1. Adiabatic Approx. \Rightarrow Drop $\frac{1}{2} \sum_i \hat{P}_i^2$

2. Ansatz: $\hat{X}_i \rightarrow x_0 + (-1)^i x_1$, where



$x_0 = -\frac{\lambda}{\omega_0^2}$ and $x_1 = \text{order parameter}$

3. Minimize the Free Energy:

$$F = \frac{N}{2} \omega_0^2 (x_0^2 + x_1^2) - \frac{2}{\beta} \sum_{\mathbf{k}} \ln(1 + e^{-\beta E_{\mathbf{k}}(x_1)})$$

4. $x_1 = 0$ corresponds to $T > T_{\text{cdw}}$

$x_1 > 0$ corresponds to $T < T_{\text{cdw}}$



Classical Monte Carlo

Monte Carlo (MC): stochastic algorithm for performing integration.

Classical Monte Carlo

Monte Carlo (MC): stochastic algorithm for performing integration.

MC in Physics: Samples Microstates at Fixed Temp \Leftrightarrow Integrates $Z = \text{Tr } e^{-\beta \hat{H}}$

Classical Monte Carlo

Monte Carlo (MC): stochastic algorithm for performing integration.

MC in Physics: Samples Microstates at Fixed Temp \Leftrightarrow Integrates $Z = \text{Tr } e^{-\beta \hat{H}}$

To Ensure Convergence to Equilibrium Distribution $\pi(\cdot)$

- **Ergodicity:** sampling protocol can access all microstates
- Irreducibility, Aperiodicity
- Detailed Balance: $\frac{\pi(A)}{\pi(B)} = \frac{P(B \rightarrow A)}{P(A \rightarrow B)} = \left[\frac{T(B \rightarrow A)}{T(A \rightarrow B)} \frac{p(B \rightarrow A)}{p(A \rightarrow B)} \right]$

Classical Monte Carlo

Monte Carlo (MC): stochastic algorithm for performing integration.

MC in Physics: Samples Microstates at Fixed Temp \Leftrightarrow Integrates $Z = \text{Tr } e^{-\beta \hat{H}}$

To Ensure Convergence to Equilibrium Distribution $\pi(\cdot)$

- **Ergodicity**: sampling protocol can access all microstates
- Irreducibility, Aperiodicity
- Detailed Balance: $\frac{\pi(A)}{\pi(B)} = \frac{P(B \rightarrow A)}{P(A \rightarrow B)} = \left[\frac{T(B \rightarrow A)}{T(A \rightarrow B)} \frac{p(B \rightarrow A)}{p(A \rightarrow B)} \right]$

Metropolis-Hastings: $p(B \rightarrow A) = \min \left[1, \frac{\pi(A)}{\pi(B)} \frac{T(A \rightarrow B)}{T(B \rightarrow A)} \right]$

Classical Monte Carlo

Monte Carlo (MC): stochastic algorithm for performing integration.

MC in Physics: Samples Microstates at Fixed Temp \Leftrightarrow Integrates $Z = \text{Tr } e^{-\beta \hat{H}}$

To Ensure Convergence to Equilibrium Distribution $\pi(\cdot)$

- **Ergodicity**: sampling protocol can access all microstates
- Irreducibility, Aperiodicity
- Detailed Balance: $\frac{\pi(A)}{\pi(B)} = \frac{P(B \rightarrow A)}{P(A \rightarrow B)} = \left[\frac{T(B \rightarrow A)}{T(A \rightarrow B)} \frac{p(B \rightarrow A)}{p(A \rightarrow B)} \right]$

Metropolis-Hastings: $p(B \rightarrow A) = \min \left[1, \frac{\pi(A)}{\pi(B)} \frac{T(A \rightarrow B)}{T(B \rightarrow A)} \right]$

$$\frac{\pi(A)}{\pi(B)} = \frac{e^{-\beta E_A}/Z}{e^{-\beta E_B}/Z} = e^{-\beta(E_A - E_B)} = e^{-\beta \Delta E}$$

Quantum Monte Carlo: Quantum Harmonic Oscillator

$$Z = \text{Tr} \left[e^{-\beta \hat{H}} \right] \quad \text{where} \quad \hat{H} = \frac{1}{2} \hat{P}^2 + \frac{\omega_0^2}{2} \hat{X}^2$$

$$Z = \text{Tr} \left[e^{-\Delta\tau \hat{H}} \dots e^{-\Delta\tau \hat{H}} \right] \quad \text{where} \quad \tau = 1 \dots L_\tau \quad \text{and} \quad \beta = L_\tau \Delta\tau$$

Suzuki-Trotter Approximation: $e^{-\Delta\tau(\hat{A}+\hat{B})} \approx e^{-\Delta\tau\hat{A}} e^{-\Delta\tau\hat{B}} + O(\Delta\tau^2)$

$$Z \approx \text{Tr} \left[e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} \dots e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} \right]$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} \langle x_1 | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} | x_2 \rangle \langle x_2 | \dots | x_{L_\tau} \rangle$$

$$\langle x_{L_\tau} | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} | x_1 \rangle$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} e^{-\Delta\tau \left(\frac{\omega_0^2}{2} \sum \hat{x}_\tau^2 \right)} \dots \int_{-\infty}^{\infty} dp \langle x_\tau | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} | p \rangle \langle p | x_{\tau+1} \rangle \dots$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} e^{-\Delta\tau \left(\frac{\omega_0^2}{2} \sum \hat{x}_\tau^2 \right)} \dots \int_{-\infty}^{\infty} e^{-\frac{\Delta\tau}{2} p^2 + ip(x_{\tau+1} - x_\tau)} \dots$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} \exp \left\{ -\Delta\tau \left[\frac{\omega_0^2}{2} \sum \hat{x}_\tau^2 + \frac{1}{2} \sum \left(\frac{x_{\tau+1} - x_\tau}{\Delta\tau} \right)^2 \right] \right\} = \int e^{-S_{\text{Lange}}(\{x\})}$$

Quantum Monte Carlo: Quantum Harmonic Oscillator

$$Z = \text{Tr} \left[e^{-\beta \hat{H}} \right] \quad \text{where} \quad \hat{H} = \frac{1}{2} \hat{P}^2 + \frac{\omega_0^2}{2} \hat{X}^2$$

$$Z = \text{Tr} \left[e^{-\Delta\tau \hat{H}} \dots e^{-\Delta\tau \hat{H}} \right] \quad \text{where} \quad \tau = 1 \dots L_\tau \quad \text{and} \quad \beta = L_\tau \Delta\tau$$

Suzuki-Trotter Approximation: $e^{-\Delta\tau(A+B)} \approx e^{-\Delta\tau\hat{A}} e^{-\Delta\tau\hat{B}} + O(\Delta\tau^2)$

$$Z \approx \text{Tr} \left[e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} \dots e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} \right]$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} \langle x_1 | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} | x_2 \rangle \langle x_2 | \dots | x_{L_\tau} \rangle$$

$$\langle x_{L_\tau} | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} | x_1 \rangle$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} e^{-\Delta\tau \left(\frac{\omega_0^2}{2} \sum \hat{x}_\tau^2 \right)} \dots \int_{-\infty}^{\infty} dp \langle x_\tau | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} | p \rangle \langle p | x_{\tau+1} \rangle \dots$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} e^{-\Delta\tau \left(\frac{\omega_0^2}{2} \sum \hat{x}_\tau^2 \right)} \dots \int_{-\infty}^{\infty} e^{-\frac{\Delta\tau}{2} p^2 + ip(x_{\tau+1} - x_\tau)} \dots$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} \exp \left\{ -\Delta\tau \left[\frac{\omega_0^2}{2} \sum \hat{x}_\tau^2 + \frac{1}{2} \sum \left(\frac{x_{\tau+1} - x_\tau}{\Delta\tau} \right)^2 \right] \right\} = \int e^{-S_{\text{Lange}}(\{x\})}$$

Quantum Monte Carlo: Quantum Harmonic Oscillator

$$Z = \text{Tr} \left[e^{-\beta \hat{H}} \right] \quad \text{where} \quad \hat{H} = \frac{1}{2} \hat{P}^2 + \frac{\omega_0^2}{2} \hat{X}^2$$

$$Z = \text{Tr} \left[e^{-\Delta\tau \hat{H}} \dots e^{-\Delta\tau \hat{H}} \right] \quad \text{where} \quad \tau = 1 \dots L_\tau \quad \text{and} \quad \beta = L_\tau \Delta\tau$$

Suzuki-Trotter Approximation: $e^{-\Delta\tau(\hat{A}+\hat{B})} \approx e^{-\Delta\tau\hat{A}}e^{-\Delta\tau\hat{B}} + \mathcal{O}(\Delta\tau^2)$

$$Z \approx \text{Tr} \left[e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} \dots e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} \right]$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} \langle x_1 | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} | x_2 \rangle \langle x_2 | \dots | x_{L_\tau} \rangle$$

$$\langle x_{L_\tau} | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} | x_1 \rangle$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} e^{-\Delta\tau \left(\frac{\omega_0^2}{2} \sum \hat{x}_\tau^2 \right)} \dots \int_{-\infty}^{\infty} dp \langle x_\tau | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} | p \rangle \langle p | x_{\tau+1} \rangle \dots$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} e^{-\Delta\tau \left(\frac{\omega_0^2}{2} \sum \hat{x}_\tau^2 \right)} \dots \int_{-\infty}^{\infty} e^{-\frac{\Delta\tau}{2} p^2 + ip(x_{\tau+1} - x_\tau)} \dots$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} \exp \left\{ -\Delta\tau \left[\frac{\omega_0^2}{2} \sum \hat{x}_\tau^2 + \frac{1}{2} \sum \left(\frac{x_{\tau+1} - x_\tau}{\Delta\tau} \right)^2 \right] \right\} = \int e^{-S_{\text{Lange}}(\{x\})}$$

Quantum Monte Carlo: Quantum Harmonic Oscillator

$$Z = \text{Tr} \left[e^{-\beta \hat{H}} \right] \quad \text{where} \quad \hat{H} = \frac{1}{2} \hat{P}^2 + \frac{\omega_0^2}{2} \hat{X}^2$$

$$Z = \text{Tr} \left[e^{-\Delta\tau \hat{H}} \dots e^{-\Delta\tau \hat{H}} \right] \quad \text{where} \quad \tau = 1 \dots L_\tau \quad \text{and} \quad \beta = L_\tau \Delta\tau$$

Suzuki-Trotter Approximation: $e^{-\Delta\tau(\hat{A}+\hat{B})} \approx e^{-\Delta\tau\hat{A}}e^{-\Delta\tau\hat{B}} + \mathcal{O}(\Delta\tau^2)$

$$Z \approx \text{Tr} \left[e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} \dots e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} \right]$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} \langle x_1 | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} | x_2 \rangle \langle x_2 | \dots | x_{L_\tau} \rangle$$

$$\langle x_{L_\tau} | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} | x_1 \rangle$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} e^{-\Delta\tau \left(\frac{\omega_0^2}{2} \sum \dot{x}_\tau^2 \right)} \dots \int_{-\infty}^{\infty} dp \langle x_\tau | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} | p \rangle \langle p | x_{\tau+1} \rangle \dots$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} e^{-\Delta\tau \left(\frac{\omega_0^2}{2} \sum \dot{x}_\tau^2 \right)} \dots \int_{-\infty}^{\infty} e^{-\frac{\Delta\tau}{2} p^2 + ip(x_{\tau+1} - x_\tau)} \dots$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} \exp \left\{ -\Delta\tau \left[\frac{\omega_0^2}{2} \sum \dot{x}_\tau^2 + \frac{1}{2} \sum \left(\frac{x_{\tau+1} - x_\tau}{\Delta\tau} \right)^2 \right] \right\} = \int e^{-S_{\text{Lange}}(x_\tau)}$$

Quantum Monte Carlo: Quantum Harmonic Oscillator

$$Z = \text{Tr} \left[e^{-\beta \hat{H}} \right] \quad \text{where} \quad \hat{H} = \frac{1}{2} \hat{P}^2 + \frac{\omega_0^2}{2} \hat{X}^2$$

$$Z = \text{Tr} \left[e^{-\Delta\tau \hat{H}} \dots e^{-\Delta\tau \hat{H}} \right] \quad \text{where} \quad \tau = 1 \dots L_\tau \quad \text{and} \quad \beta = L_\tau \Delta\tau$$

Suzuki-Trotter Approximation: $e^{-\Delta\tau(\hat{A}+\hat{B})} \approx e^{-\Delta\tau\hat{A}} e^{-\Delta\tau\hat{B}} + \mathcal{O}(\Delta\tau^2)$

$$Z \approx \text{Tr} \left[e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} \dots e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} \right]$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} \langle x_1 | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} | x_2 \rangle \langle x_2 | \dots | x_{L_\tau} \rangle$$

$$\langle x_{L_\tau} | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} | x_1 \rangle$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} e^{-\Delta\tau \left(\frac{\omega_0^2}{2} \sum \hat{x}_\tau^2 \right)} \dots \int_{-\infty}^{\infty} dp \langle x_\tau | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} | p \rangle \langle p | x_{\tau+1} \rangle \dots$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} e^{-\Delta\tau \left(\frac{\omega_0^2}{2} \sum \hat{x}_\tau^2 \right)} \dots \int_{-\infty}^{\infty} e^{-\frac{\Delta\tau}{2} p^2 + ip(x_{\tau+1} - x_\tau)} \dots$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} \exp \left\{ -\Delta\tau \left[\frac{\omega_0^2}{2} \sum \hat{x}_\tau^2 + \frac{1}{2} \sum \left(\frac{x_{\tau+1} - x_\tau}{\Delta\tau} \right)^2 \right] \right\} = \int e^{-S_{\text{Lange}}(x_\tau)}$$

Quantum Monte Carlo: Quantum Harmonic Oscillator

$$Z = \text{Tr} \left[e^{-\beta \hat{H}} \right] \quad \text{where} \quad \hat{H} = \frac{1}{2} \hat{P}^2 + \frac{\omega_0^2}{2} \hat{X}^2$$

$$Z = \text{Tr} \left[e^{-\Delta\tau \hat{H}} \dots e^{-\Delta\tau \hat{H}} \right] \quad \text{where} \quad \tau = 1 \dots L_\tau \quad \text{and} \quad \beta = L_\tau \Delta\tau$$

Suzuki-Trotter Approximation: $e^{-\Delta\tau(\hat{A}+\hat{B})} \approx e^{-\Delta\tau\hat{A}} e^{-\Delta\tau\hat{B}} + \mathcal{O}(\Delta\tau^2)$

$$Z \approx \text{Tr} \left[e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} \dots e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} \right]$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} \langle x_1 | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} | x_2 \rangle \langle x_2 | \dots | x_{L_\tau} \rangle$$

$$\langle x_{L_\tau} | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} | x_1 \rangle$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} e^{-\Delta\tau \left(\frac{\omega_0^2}{2} \sum_{\tau} x_{\tau}^2 \right)} \dots \int_{-\infty}^{\infty} dp \langle x_{\tau} | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} | p \rangle \langle p | x_{\tau+1} \rangle \dots$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} e^{-\Delta\tau \left(\frac{\omega_0^2}{2} \sum_{\tau} x_{\tau}^2 \right)} \dots \int_{-\infty}^{\infty} e^{-\frac{\Delta\tau}{2} p^2 + ip(x_{\tau+1} - x_{\tau})} \dots$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} \exp \left\{ -\Delta\tau \left[\frac{\omega_0^2}{2} \sum_{\tau} x_{\tau}^2 + \frac{1}{2} \sum_{\tau} \left(\frac{x_{\tau+1} - x_{\tau}}{\Delta\tau} \right)^2 \right] \right\} = \int e^{-S_{\text{Langevin}}(x_{\tau})}$$

Quantum Monte Carlo: Quantum Harmonic Oscillator

$$Z = \text{Tr} \left[e^{-\beta \hat{H}} \right] \quad \text{where} \quad \hat{H} = \frac{1}{2} \hat{P}^2 + \frac{\omega_0^2}{2} \hat{X}^2$$

$$Z = \text{Tr} \left[e^{-\Delta\tau \hat{H}} \dots e^{-\Delta\tau \hat{H}} \right] \quad \text{where} \quad \tau = 1 \dots L_\tau \quad \text{and} \quad \beta = L_\tau \Delta\tau$$

Suzuki-Trotter Approximation: $e^{-\Delta\tau(\hat{A}+\hat{B})} \approx e^{-\Delta\tau\hat{A}} e^{-\Delta\tau\hat{B}} + \mathcal{O}(\Delta\tau^2)$

$$Z \approx \text{Tr} \left[e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} \dots e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} \right]$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} \langle x_1 | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} | x_2 \rangle \langle x_2 | \dots | x_{L_\tau} \rangle$$

$$\langle x_{L_\tau} | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} | x_1 \rangle$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} e^{-\Delta\tau \left(\frac{\omega_0^2}{2} \sum_{\tau} x_{\tau}^2 \right)} \dots \int_{-\infty}^{\infty} dp \langle x_{\tau} | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} | p \rangle \langle p | x_{\tau+1} \rangle \dots$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} e^{-\Delta\tau \left(\frac{\omega_0^2}{2} \sum_{\tau} x_{\tau}^2 \right)} \dots \int_{-\infty}^{\infty} e^{-\frac{\Delta\tau}{2} p^2 + ip(x_{\tau+1} - x_{\tau})} \dots$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} \exp \left\{ -\Delta\tau \left[\frac{1}{2} \sum_{\tau} x_{\tau}^2 + \frac{1}{2} \sum_{\tau} \left(\frac{x_{\tau+1} - x_{\tau}}{\Delta\tau} \right)^2 \right] \right\} = \int e^{-S_{\text{classical}}(x)}$$

Quantum Monte Carlo: Quantum Harmonic Oscillator

$$Z = \text{Tr} \left[e^{-\beta \hat{H}} \right] \quad \text{where} \quad \hat{H} = \frac{1}{2} \hat{P}^2 + \frac{\omega_0^2}{2} \hat{X}^2$$

$$Z = \text{Tr} \left[e^{-\Delta\tau \hat{H}} \dots e^{-\Delta\tau \hat{H}} \right] \quad \text{where} \quad \tau = 1 \dots L_\tau \quad \text{and} \quad \beta = L_\tau \Delta\tau$$

Suzuki-Trotter Approximation: $e^{-\Delta\tau(\hat{A}+\hat{B})} \approx e^{-\Delta\tau\hat{A}} e^{-\Delta\tau\hat{B}} + \mathcal{O}(\Delta\tau^2)$

$$Z \approx \text{Tr} \left[e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} \dots e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} \right]$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} \langle x_1 | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} | x_2 \rangle \langle x_2 | \dots | x_{L_\tau} \rangle$$

$$\langle x_{L_\tau} | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} e^{-\Delta\tau \left(\frac{1}{2} \omega_0^2 \hat{X}^2 \right)} | x_1 \rangle$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} e^{-\Delta\tau \left(\frac{\omega_0^2}{2} \sum_{\tau} x_{\tau}^2 \right)} \dots \int_{-\infty}^{\infty} dp \langle x_{\tau} | e^{-\Delta\tau \left(\frac{1}{2} \hat{P}^2 \right)} | p \rangle \langle p | x_{\tau+1} \rangle \dots$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} e^{-\Delta\tau \left(\frac{\omega_0^2}{2} \sum_{\tau} x_{\tau}^2 \right)} \dots \int_{-\infty}^{\infty} e^{-\frac{\Delta\tau}{2} p^2 + ip(x_{\tau+1} - x_{\tau})} \dots$$

$$Z = \int_{-\infty}^{\infty} dx_1 \dots dx_{L_\tau} \exp \left\{ -\Delta\tau \left[\frac{\omega_0^2}{2} \sum_{\tau} x_{\tau}^2 + \frac{1}{2} \sum_{\tau} \left(\frac{x_{\tau+1} - x_{\tau}}{\Delta\tau} \right)^2 \right] \right\} = \int e^{-S_{\text{bose}}(\{x_{\tau}\})}$$

Determinant Quantum Monte Carlo

$$Z = \text{Tr} \left[e^{-\Delta\tau(\hat{U} + \hat{K} + \hat{V})} \dots e^{-\Delta\tau(\hat{U} + \hat{K} + \hat{V})} \right]$$

$$Z \approx \text{Tr} \left[e^{-\Delta\tau\hat{U}} e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}} \dots e^{-\Delta\tau\hat{U}} e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}} \right]$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,j} dx_{\tau,j} e^{-S_{\text{bond}}(\{x_{\tau,j}\})} \prod_{\sigma} \text{Tr} \left[e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}(1)} \dots e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}(L_\tau)} \right]$$

Note: $\hat{V}(\tau) = \hat{c}^\dagger \tilde{V}(\tau) \hat{c}$ and $\hat{K} = \hat{c}^\dagger \tilde{K} \hat{c}$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,j} dx_{\tau,j} e^{-S_{\text{bond}}(\{x_{\tau,j}\})} \prod_{\sigma} \det \left[1 + e^{-\Delta\tau\tilde{K}} e^{-\Delta\tau\tilde{V}(1)} \dots e^{-\Delta\tau\tilde{K}} e^{-\Delta\tau\tilde{V}(L_\tau)} \right]$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,j} dx_{\tau,j} e^{-S_{\text{bond}}(\{x_{\tau,j}\})} \det [\tilde{M}_\uparrow(\{x_{\tau,j}\})] \det [\tilde{M}_\downarrow(\{x_{\tau,j}\})]$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,j} dx_{\tau,j} \left\{ e^{-S_{\text{bond}}(\{x_{\tau,j}\})} \det [\tilde{M}(\{x_{\tau,j}\})] \right\}^2$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,j} dx_{\tau,j} W(\{x_{\tau,j}\})$$

Determinant Quantum Monte Carlo

$$Z = \text{Tr} \left[e^{-\Delta\tau(\hat{U} + \hat{K} + \hat{V})} \dots e^{-\Delta\tau(\hat{U} + \hat{K} + \hat{V})} \right]$$

$$Z \approx \text{Tr} \left[e^{-\Delta\tau\hat{U}} e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}} \dots e^{-\Delta\tau\hat{U}} e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}} \right]$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,i} dx_{\tau,i} e^{-S_{\text{bose}}(\{x_{\tau,i}\})} \prod_{\sigma} \text{Tr} \left[e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}(1)} \dots e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}(L_\tau)} \right]$$

Note: $\hat{V}(\tau) = \hat{c}^\dagger \tilde{V}(\tau) \hat{c}$ and $\hat{K} = \hat{c}^\dagger \tilde{K} \hat{c}$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,j} dx_{\tau,j} e^{-S_{\text{bose}}(\{x_{\tau,j}\})} \prod_{\sigma} \det \left[1 + e^{-\Delta\tau\tilde{K}} e^{-\Delta\tau\tilde{V}(1)} \dots e^{-\Delta\tau\tilde{K}} e^{-\Delta\tau\tilde{V}(L_\tau)} \right]$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,j} dx_{\tau,j} e^{-S_{\text{bose}}(\{x_{\tau,j}\})} \det [\tilde{M}_\uparrow(\{x_{\tau,j}\})] \det [\tilde{M}_\downarrow(\{x_{\tau,j}\})]$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,j} dx_{\tau,j} \left\{ e^{-S_{\text{bose}}(\{x_{\tau,j}\})} \det [\tilde{M}(\{x_{\tau,j}\})] \right\}^2$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,j} dx_{\tau,j} W(\{x_{\tau,j}\})$$

Determinant Quantum Monte Carlo

$$Z = \text{Tr} \left[e^{-\Delta\tau(\hat{U} + \hat{K} + \hat{V})} \dots e^{-\Delta\tau(\hat{U} + \hat{K} + \hat{V})} \right]$$

$$Z \approx \text{Tr} \left[e^{-\Delta\tau\hat{U}} e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}} \dots e^{-\Delta\tau\hat{U}} e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}} \right]$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,i} dx_{\tau,i} e^{-S_{\text{bose}}(\{x_{\tau,i}\})} \prod_{\sigma} \text{Tr} \left[e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}(1)} \dots e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}(L_\tau)} \right]$$

Note: $\hat{V}(\tau) = \hat{\mathbf{c}}^\dagger \bar{V}(\tau) \hat{\mathbf{c}}$ and $\hat{K} = \hat{\mathbf{c}}^\dagger \bar{K} \hat{\mathbf{c}}$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,j} dx_{\tau,j} e^{-S_{\text{bose}}(\{x_{\tau,j}\})} \prod_{\sigma} \det \left[1 + e^{-\Delta\tau\bar{K}} e^{-\Delta\tau\bar{V}(1)} \dots e^{-\Delta\tau\bar{K}} e^{-\Delta\tau\bar{V}(L_\tau)} \right]$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,j} dx_{\tau,j} e^{-S_{\text{bose}}(\{x_{\tau,j}\})} \det [\bar{M}_\uparrow(\{x_{\tau,j}\})] \det [\bar{M}_\downarrow(\{x_{\tau,j}\})]$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,j} dx_{\tau,j} \left\{ e^{-S_{\text{bose}}(\{x_{\tau,j}\})} \det [\bar{M}(\{x_{\tau,j}\})] \right\}^2$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,j} dx_{\tau,j} W(\{x_{\tau,j}\})$$

Determinant Quantum Monte Carlo

$$Z = \text{Tr} \left[e^{-\Delta\tau(\hat{U} + \hat{K} + \hat{V})} \dots e^{-\Delta\tau(\hat{U} + \hat{K} + \hat{V})} \right]$$

$$Z \approx \text{Tr} \left[e^{-\Delta\tau\hat{U}} e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}} \dots e^{-\Delta\tau\hat{U}} e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}} \right]$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,i} dx_{\tau,i} e^{-S_{\text{bose}}(\{x_{\tau,i}\})} \prod_{\sigma} \text{Tr} \left[e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}(1)} \dots e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}(L_\tau)} \right]$$

Note: $\hat{V}(\tau) = \hat{\mathbf{c}}^\dagger \bar{V}(\tau) \hat{\mathbf{c}}$ and $\hat{K} = \hat{\mathbf{c}}^\dagger \bar{K} \hat{\mathbf{c}}$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,i} dx_{\tau,i} e^{-S_{\text{bose}}(\{x_{\tau,i}\})} \prod_{\sigma} \det \left[\mathbb{1} + e^{-\Delta\tau\bar{K}} e^{-\Delta\tau\bar{V}(1)} \dots e^{-\Delta\tau\bar{K}} e^{-\Delta\tau\bar{V}(L_\tau)} \right]$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,j} dx_{\tau,j} e^{-S_{\text{bose}}(\{x_{\tau,j}\})} \det [\bar{M}_\uparrow(\{x_{\tau,j}\})] \det [\bar{M}_\downarrow(\{x_{\tau,j}\})]$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,j} dx_{\tau,j} \left\{ e^{-S_{\text{bose}}(\{x_{\tau,j}\})} \det [\bar{M}(\{x_{\tau,j}\})] \right\}^2$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,j} dx_{\tau,j} W(\{x_{\tau,j}\})$$

Determinant Quantum Monte Carlo

$$Z = \text{Tr} \left[e^{-\Delta\tau(\hat{U} + \hat{K} + \hat{V})} \dots e^{-\Delta\tau(\hat{U} + \hat{K} + \hat{V})} \right]$$

$$Z \approx \text{Tr} \left[e^{-\Delta\tau\hat{U}} e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}} \dots e^{-\Delta\tau\hat{U}} e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}} \right]$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,i} dx_{\tau,i} e^{-S_{\text{bose}}(\{x_{\tau,i}\})} \prod_{\sigma} \text{Tr} \left[e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}(1)} \dots e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}(L_\tau)} \right]$$

Note: $\hat{V}(\tau) = \hat{\mathbf{c}}^\dagger \bar{V}(\tau) \hat{\mathbf{c}}$ and $\hat{K} = \hat{\mathbf{c}}^\dagger \bar{K} \hat{\mathbf{c}}$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,i} dx_{\tau,i} e^{-S_{\text{bose}}(\{x_{\tau,i}\})} \prod_{\sigma} \det \left[\mathbb{1} + e^{-\Delta\tau\bar{K}} e^{-\Delta\tau\bar{V}(1)} \dots e^{-\Delta\tau\bar{K}} e^{-\Delta\tau\bar{V}(L_\tau)} \right]$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,i} dx_{\tau,i} e^{-S_{\text{bose}}(\{x_{\tau,i}\})} \det [\bar{M}_\uparrow(\{x_{\tau,i}\})] \det [\bar{M}_\downarrow(\{x_{\tau,i}\})]$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,i} dx_{\tau,i} \left\{ e^{-S_{\text{bose}}(\{x_{\tau,i}\})} \det [\bar{M}(\{x_{\tau,i}\})]^2 \right\}$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,i} dx_{\tau,i} W(\{x_{\tau,i}\})$$

Determinant Quantum Monte Carlo

$$Z = \text{Tr} \left[e^{-\Delta\tau(\hat{U} + \hat{K} + \hat{V})} \dots e^{-\Delta\tau(\hat{U} + \hat{K} + \hat{V})} \right]$$

$$Z \approx \text{Tr} \left[e^{-\Delta\tau\hat{U}} e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}} \dots e^{-\Delta\tau\hat{U}} e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}} \right]$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,i} dx_{\tau,i} e^{-S_{\text{bose}}(\{x_{\tau,i}\})} \prod_{\sigma} \text{Tr} \left[e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}(1)} \dots e^{-\Delta\tau\hat{K}} e^{-\Delta\tau\hat{V}(L_{\tau})} \right]$$

Note: $\hat{V}(\tau) = \hat{\mathbf{c}}^\dagger \bar{V}(\tau) \hat{\mathbf{c}}$ and $\hat{K} = \hat{\mathbf{c}}^\dagger \bar{K} \hat{\mathbf{c}}$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,i} dx_{\tau,i} e^{-S_{\text{bose}}(\{x_{\tau,i}\})} \prod_{\sigma} \det \left[1 + e^{-\Delta\tau\bar{K}} e^{-\Delta\tau\bar{V}(1)} \dots e^{-\Delta\tau\bar{K}} e^{-\Delta\tau\bar{V}(L_{\tau})} \right]$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,i} dx_{\tau,i} e^{-S_{\text{bose}}(\{x_{\tau,i}\})} \det [\bar{M}_{\uparrow}(\{x_{\tau,i}\})] \det [\bar{M}_{\downarrow}(\{x_{\tau,i}\})]$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,i} dx_{\tau,i} \left\{ e^{-S_{\text{bose}}(\{x_{\tau,i}\})} \det [\bar{M}(\{x_{\tau,i}\})]^2 \right\}$$

$$Z = \int_{-\infty}^{\infty} \prod_{\tau,i} dx_{\tau,i} W(\{x_{\tau,i}\})$$

DQMC Measurements

$$G_{i,j} = \langle c_i c_j^\dagger \rangle = \left[\mathbb{1} + e^{-\Delta\tau \bar{K}} e^{-\Delta\tau \bar{V}(1)} \dots e^{-\Delta\tau \bar{K}} e^{-\Delta\tau \bar{V}(L_\tau)} \right]^{-1} = \bar{M}_{i,j}^{-1}$$

DQMC Measurements

$$G_{i,j} = \langle c_i c_j^\dagger \rangle = \left[\mathbb{1} + e^{-\Delta\tau \bar{K}} e^{-\Delta\tau \bar{V}(1)} \dots e^{-\Delta\tau \bar{K}} e^{-\Delta\tau \bar{V}(L_\tau)} \right]^{-1} = \bar{M}_{i,j}^{-1}$$

Detecting CDW Order:

- Density-Density Correlation: $c(\mathbf{r}) = \langle n_{\mathbf{i}} n_{\mathbf{i+r}} \rangle = \langle (n_{\uparrow,\mathbf{i}} + n_{\downarrow,\mathbf{i}})(n_{\uparrow,\mathbf{i+r}} + n_{\downarrow,\mathbf{i+r}}) \rangle$

DQMC Measurements

$$G_{i,j} = \langle c_i c_j^\dagger \rangle = \left[\mathbb{1} + e^{-\Delta\tau \tilde{K}} e^{-\Delta\tau \tilde{V}(1)} \dots e^{-\Delta\tau \tilde{K}} e^{-\Delta\tau \tilde{V}(L_\tau)} \right]^{-1} = \bar{M}_{i,j}^{-1}$$

Detecting CDW Order:


- Density-Density Correlation: $c(\mathbf{r}) = \langle n_{\mathbf{i}} n_{\mathbf{i+r}} \rangle = \langle (n_{\uparrow,\mathbf{i}} + n_{\downarrow,\mathbf{i}})(n_{\uparrow,\mathbf{i+r}} + n_{\downarrow,\mathbf{i+r}}) \rangle$
- CDW Structure Factor: $S_{\text{cdw}} = S(\mathbf{q}_{\text{cdw}}) = \sum_{n=1}^N e^{i\mathbf{q}_{\text{cdw}} \cdot \mathbf{r}_n} c(\mathbf{r}_n) = \sum_{n=1}^N (-1)^n c(\mathbf{r}_n)$

DQMC Measurements

$$G_{i,j} = \langle c_i c_j^\dagger \rangle = \left[\mathbb{1} + e^{-\Delta\tau \bar{K}} e^{-\Delta\tau \bar{V}(1)} \dots e^{-\Delta\tau \bar{K}} e^{-\Delta\tau \bar{V}(L_\tau)} \right]^{-1} = \bar{M}_{i,j}^{-1}$$

Detecting CDW Order:

- Density-Density Correlation: $c(\mathbf{r}) = \langle n_{\mathbf{i}} n_{\mathbf{i}+\mathbf{r}} \rangle = \langle (n_{\uparrow,\mathbf{i}} + n_{\downarrow,\mathbf{i}})(n_{\uparrow,\mathbf{i}+\mathbf{r}} + n_{\downarrow,\mathbf{i}+\mathbf{r}}) \rangle$
- CDW Structure Factor: $S_{\text{cdw}} = S(\mathbf{q}_{\text{cdw}}) = \sum_{n=1}^N e^{i\mathbf{q}_{\text{cdw}} \cdot \mathbf{r}_n} c(\mathbf{r}_n) = \sum_{n=1}^N (-1)^n c(\mathbf{r}_n)$



- $T > T_{\text{cdw}} \implies S_{\text{cdw}} \perp N$
- $T < T_{\text{cdw}} \implies S_{\text{cdw}} \sim N$
- Perfect CDW $\implies S_{\text{cdw}} = N$

Finite Size Scaling: Extracting T_c

1. Consider a singular quantity: $Q \sim |t|^{-\kappa}$ where $t = \frac{T-T_c}{T_c}$

Finite Size Scaling: Extracting T_c

1. Consider a singular quantity: $Q \sim |t|^{-\kappa}$ where $t = \frac{T-T_c}{T_c}$
2. $\xi \sim |t|^{-\nu} \implies |t| \sim \xi^{-1/\nu}$

Finite Size Scaling: Extracting T_c

1. Consider a singular quantity: $Q \sim |t|^{-\kappa}$ where $t = \frac{T-T_c}{T_c}$
2. $\xi \sim |t|^{-\nu} \implies |t| \sim \xi^{-1/\nu}$
3. $Q \sim \xi^{\kappa/\nu} \implies Q_{\max} \sim L^{\kappa/\nu}$ (Finite Size Effects when $\xi \approx L$)

Finite Size Scaling: Extracting T_c

1. Consider a singular quantity: $Q \sim |t|^{-\kappa}$ where $t = \frac{T-T_c}{T_c}$
2. $\xi \sim |t|^{-\nu} \implies |t| \sim \xi^{-1/\nu}$
3. $Q \sim \xi^{\kappa/\nu} \implies Q_{\max} \sim L^{\kappa/\nu}$ (Finite Size Effects when $\xi \approx L$)
4. $|t_{\max}(L)| \sim L^{-1/\nu} \implies$ shifted T_c because L is finite

Finite Size Scaling: Extracting T_c

1. Consider a singular quantity: $Q \sim |t|^{-\kappa}$ where $t = \frac{T-T_c}{T_c}$
2. $\xi \sim |t|^{-\nu} \implies |t| \sim \xi^{-1/\nu}$
3. $Q \sim \xi^{\kappa/\nu} \implies Q_{\max} \sim L^{\kappa/\nu}$ (Finite Size Effects when $\xi \approx L$)
4. $|t_{\max}(L)| \sim L^{-1/\nu} \implies$ shifted T_c because L is finite
5. Finite-Size Scaling Hypothesis:

$$Q(t, L) = L^\sigma f(\xi/L) = L^\sigma g'(|t|^{-\nu}/L) = L^\sigma g(t L^{-1/\nu})$$

Finite Size Scaling: Extracting T_c

1. Consider a singular quantity: $Q \sim |t|^{-\kappa}$ where $t = \frac{T-T_c}{T_c}$
2. $\xi \sim |t|^{-\nu} \implies |t| \sim \xi^{-1/\nu}$
3. $Q \sim \xi^{\kappa/\nu} \implies Q_{\max} \sim L^{\kappa/\nu}$ (Finite Size Effects when $\xi \approx L$)
4. $|t_{\max}(L)| \sim L^{-1/\nu} \implies$ shifted T_c because L is finite
5. Finite-Size Scaling Hypothesis:

$$Q(t, L) = L^\sigma f(\xi/L) = L^\sigma g'(|t|^{-\nu}/L) = L^\sigma g(t L^{-1/\nu})$$

6. $L \rightarrow \infty \implies Q \sim |t|^{-\kappa} \implies g(x) \sim x^{-\kappa} \implies Q = L^{\kappa/\nu} g(t L^{-1/\nu})$

Scaling Behavior of S_{cdw}

1. Order Parameter: $\psi = \langle n_A - n_B \rangle \sim |t|^\beta$

Scaling Behavior of S_{cdw}

1. Order Parameter: $\psi = \langle n_A - n_B \rangle \sim |t|^\beta$
2. $[c(\mathbf{r}) = \langle n_{\mathbf{i}} n_{\mathbf{i}+\mathbf{r}} \rangle] \sim \psi^2 \implies c(\mathbf{r}) \sim |t|^{2\beta}$

Scaling Behavior of S_{cdw}

1. Order Parameter: $\psi = \langle n_A - n_B \rangle \sim |t|^\beta$
2. $[c(\mathbf{r}) = \langle n_{\mathbf{i}} n_{\mathbf{i}+\mathbf{r}} \rangle] \sim \psi^2 \implies c(\mathbf{r}) \sim |t|^{2\beta}$
3. $S_{\text{cdw}} \sim L^2 c(\mathbf{r}) \implies S_{\text{cdw}} \sim L^2 |t|^{2\beta}$

Scaling Behavior of S_{cdw}


1. Order Parameter: $\psi = \langle n_A - n_B \rangle \sim |t|^\beta$
2. $[c(\mathbf{r}) = \langle n_{\mathbf{i}} n_{\mathbf{i}+\mathbf{r}} \rangle] \sim \psi^2 \implies c(\mathbf{r}) \sim |t|^{2\beta}$
3. $S_{\text{cdw}} \sim L^2 c(\mathbf{r}) \implies S_{\text{cdw}} \sim L^2 |t|^{2\beta}$
4. $L^{-2} S_{\text{cdw}} \sim |t|^{2\beta}$

Scaling Behavior of S_{cdw}

1. Order Parameter: $\psi = \langle n_A - n_B \rangle \sim |t|^\beta$
2. $[c(\mathbf{r}) = \langle n_{\mathbf{i}} n_{\mathbf{i}+\mathbf{r}} \rangle] \sim \psi^2 \implies c(\mathbf{r}) \sim |t|^{2\beta}$
3. $S_{\text{cdw}} \sim L^2 c(\mathbf{r}) \implies S_{\text{cdw}} \sim L^2 |t|^{2\beta}$
4. $L^{-2} S_{\text{cdw}} \sim |t|^{2\beta}$
5. $L^{-2} S_{\text{cdw}} = L^{-2\beta/\nu} g(t L^{-1/\nu}) \implies \boxed{L^{2\beta/\nu - 2} S_{\text{cdw}} = g(t L^{-1/\nu})}$



Getting T_{cdw} from S_{cdw} : Finite Size Scaling

Ising Universality Class: $\beta = 1/8, \nu = 1/4, \alpha = 0, \gamma = 7/4, \eta = 1/4$



Getting T_{cdw} from S_{cdw} : Finite Size Scaling


Ising Universality Class: $\beta = 1/8, \nu = 1/4, \alpha = 0, \gamma = 7/4, \eta = 1/4$



Getting T_{cdw} from S_{cdw} : Finite Size Scaling



Ising Universality Class:

$$\boxed{\beta = 1/8, \quad \nu = 1/4, \quad \alpha = 0, \quad \gamma = 7/4, \quad \eta = 1/4}$$



More Results: The Effect of δ on S_{cdw}

S_{cdw} Sensitive to δ





More Results: CDW as $\delta \rightarrow 1$

$$T_{\text{cdw}} > 0 \quad \forall \quad \delta \in [0, 1)$$



More Results: CDW as $\delta \rightarrow 1$

$$T_{\text{cdw}} > 0 \quad \forall \quad \delta \in [0, 1)$$




More Results: CDW as $\delta \rightarrow 1$

$$T_{\text{cdw}} > 0 \quad \forall \quad \delta \in [0, 1)$$



$$G(\mathbf{k}, \tau) = \int d\omega \frac{A(\mathbf{k}, \omega)}{e^{-\beta\omega} + 1} e^{-\omega\tau}$$

$$N(\omega) = \sum_{\mathbf{k}} A(\mathbf{k}, \omega)$$



Summary

- $\delta \lesssim 0.4 \implies T_{\text{cdw}}$ Insensitive to δ
- δ Suppresses S_{cdw}
- $T_{\text{cdw}} > 0 \quad \forall \quad \delta \in [0, 1)$

Future Work

- Holstein Model on Cubic Lattice using SLMC + Langevin
(ongoing)
- Move away from half-filling: striped CDW and/or SC?
- Kagome Lattice at Half-filling
- etc.

Thank You!