

MPCS 58020 2017: Homework 3 Solutions

April 30, 2017

1. **Shumway 3.3** Using eq 3.11 from Shumway:

$$\begin{aligned}
 x_t &= -\sum_{j=1}^{\infty} \phi^{-j} w_{t+j} \\
 E(x_t) &= E\left(-\sum_{j=1}^{\infty} \phi^{-j} w_{t+j}\right) \\
 &= -\sum_{j=1}^{\infty} \phi^{-j} E(w_{t+j}) \\
 &= -\sum_{j=1}^{\infty} \phi^{-j} 0 \\
 &= 0
 \end{aligned}$$

so the process is stationary with zero mean.

For the autocovariance:

$$\begin{aligned}
 \gamma(h) &= \text{cov}(x_{t+h}, x_t) = E[(x_{t+h} - \bar{x})(x_t - \bar{x})] = E[x_{t+h}x_t] \\
 &= E\left[\left(-\sum_{j=1}^{\infty} \phi^{-j} w_{t+h+j}\right)\left(-\sum_{k=1}^{\infty} \phi^{-k} w_{t+k}\right)\right] \\
 &= E[(\phi^{-1}w_{t+h+1} + \phi^{-2}w_{t+h+2} + \dots)(\phi^{-1}w_{t+1} + \dots + \phi^{-h-1}w_{t+h+1} + \phi^{-h-2}w_{t+h+2} + \dots)] \\
 &= E[\phi^{-2}w_{t+h+1}w_{t+1} + \dots + \phi^{-h-2}w_{t+h+1}^2 + \phi^{-h-4}w_{t+h+2}^2 + \dots] \\
 &= E\left[\phi^{-2}w_{t+h+1}w_{t+1} + \dots + \sum_{j=1}^{\infty} \phi^{-h-2j}w_{t+h+j}^2\right]
 \end{aligned}$$

when taking the expectation, all the terms with $w_i w_j$, $i \neq j$ will become zero, since the w_i are iid. This leaves just the sum:

$$\begin{aligned}
 \gamma(h) &= E\left[\sum_{j=1}^{\infty} \phi^{-h-2j}w_{t+h+j}^2\right] \\
 &= \phi^{-h} E\left[\sum_{j=1}^{\infty} \phi^{-2j}w_{t+h+j}^2\right] \\
 &= \phi^{-h} \sum_{j=1}^{\infty} \phi^{-2j} E(w_{t+h+j}^2) \\
 &= \sigma_w^2 \phi^{-h} \sum_{j=1}^{\infty} \phi^{-2j} \\
 &= \sigma_w^2 \phi^{-h} \phi^{-2} \sum_{j=0}^{\infty} \phi^{-2j} \\
 &= \frac{\sigma_w^2 \phi^{-h} \phi^{-2}}{1 - \phi^{-2}}
 \end{aligned}$$

Consider the series $y_t = \phi^{-1}y_{t-1} + v_t$ with $v_t \sim \text{iid } N(0, \sigma_w^2 \phi^{-2})$. Since $|\phi| > 1$, $|\phi^{-1}| < 1$, so y_t is actually a causal AR(1) process with parameters $\phi' = \phi^{-1}$ and is stationary with zero mean. Note

that the standard deviation of v_t is $\sigma_w \phi^{-1}$, so by eq 3.7, the autocovariance is:

$$\begin{aligned}\gamma(h) &= \frac{(\sigma_w \phi^{-1})^2 (\phi^{-1})^h}{1 - (\phi^{-1})^2} \\ &= \frac{\sigma_w^2 \phi^{-2} \phi^{-h}}{1 - \phi^{-2}} \\ &= \frac{\sigma_w^2 \phi^{-h} \phi^{-2}}{1 - \phi^{-2}}\end{aligned}$$

2. Shumway 3.4

(a)

$$\begin{aligned}x_t &= 0.80x_{t-1} - 0.15x_{t-2} + w_t - 0.30w_{t-1} \\ (1 - 0.80B + 0.15B^2)x_t &= (1 - 0.30B)w_t \\ (1 - 0.30B)(1 - 0.5B)x_t &= (1 - 0.30B)w_t \\ (1 - 0.5B)x_t &= w_t\end{aligned}$$

This is actually an AR(1) process with $\phi = 0.5$, because of parameter redundancy. It is causal because $|\phi| < 1$.

(b)

$$\begin{aligned}x_t &= x_{t-1} - 0.50x_{t-2} + w_t - w_{t-1} \\ (1 - B + 0.50B^2)x_t &= (1 - B)w_t\end{aligned}$$

This is an ARMA(2,1) process. Because the root of $\Theta(z) = (1 - z)$ is 1, this is not an invertible process. The roots of $\Phi(z) = 1 - z + 0.5z^2$ are $1 \pm i$, which are both outside the unit circle, so the process is causal.

3. **Shumway 3.6** For the AR(2) model $x_t = -0.9x_{t-2} + w_t$, the autoregressive polynomial is $\Phi(z) = 1 + 0.9z^2$. The roots are $\pm \frac{\sqrt{10}}{3}i$, both of which are outside the unit circle, so the model is causal. By example 3.9 (iii), the ACF has the form $\rho(h) = a|z_1|^{-h} \cos(h\theta + b)$. z_1 is one of the roots so $|z_1| = \frac{\sqrt{10}}{3}$, and θ is defined as the angle of the polar coordinate for $z_1 = |z_1|e^{i\theta}$. In this case, there is no real part, so $\theta = \pi/2$. a and b are determined based on $\rho(0) = 1$ and $\rho(1) = \phi_1/(1 - \phi_2) = 0$, since $\phi_1 = 0$. This yields $\rho(0) = 1 = a \cos(b)$ and $\rho(1) = 0 = \frac{3a}{\sqrt{10}} \cos(\pi/2 + b)$. The second equation means $b = 0$, since $\cos(\pi/2) = 0$. Plugging $b = 0$ into the first equation we get $a = 1$, since $\cos(0) = 1$. Therefore the ACF is:

$$\gamma(h) = \left(\frac{\sqrt{10}}{3} \right)^{-h} \cos\left(\frac{\pi}{2}h\right)$$