

# Report

Zihan Lin

## Motivation

Extreme weather is on the rise due to climate change, with many, especially those who come from a lower socioeconomic background, already feeling the adverse effects changing weather patterns. Asthma is a co-morbidity for other chronic diseases and also disproportionately affects those from a lower economic status. In this project, we want to focus on asthma prevalence/incidence as hotter temperatures can lead to more pollen, air pollution, and other lung irritants. Studying these trends will help understand the health risks in vulnerable populations.

Key Questions: 1. How do asthma cases correlate with temperature extremes? 2. Are there temporal or spatial patterns in asthma prevalence?

## Data

To retrieve the temperature data, different options were explored from within the `rnoaa` package. `meteo_pull_monitors()` was first explored, but due to the number of monitors from each state, it was not realistic to pull data from each monitor to calculate the average temperature over a 10-year period. `ncdc()` was then investigated for use. This uses the National Climatic Data Center to access weather data and already has a `TAVG` variable for the average temperature. However, there were limitations as it could only gather data one year at a time and there was also issues with the API of this site. Using online resources was not realistic as datasets online were typically separated by year/month/day so getting a seasonal average was not possible. In the end, the temperature data was extracted using `meteo_pull_monitors()`. 30 monitors from each state were randomly selected and the daily mean of each station was found by averaging the minimum and maximum temperatures (the `TAVG` as reported by `meteo_pull_monitors()` could not be used as many stations were missing this data) for every day from 1/1/2011 to 12/31/2021 in every state. The data was then grouped by state, year, and season to find the seasonal average for every state in every year.

```
# Load datasets
asthma_data <- read_csv("data/asthma_data.csv")

## Rows: 559 Columns: 3
## -- Column specification -----
## Delimiter: ","
## chr (1): state
## dbl (2): year_name, prevalence_percent
##
## i Use 'spec()' to retrieve the full column specification for this data.
## i Specify the column types or set 'show_col_types = FALSE' to quiet this message.

temp_data <- read_csv("data/temp_data.csv")

## Rows: 2193 Columns: 4
```

```
## -- Column specification -----
## Delimiter: ","
## chr (2): state, season
## dbl (2): year, avg_temp
##
## i Use 'spec()' to retrieve the full column specification for this data.
## i Specify the column types or set 'show_col_types = FALSE' to quiet this message.
```

The analysis uses two primary datasets: Asthma Data and Temperature Data. The Asthma Data consists of 559 observations and 3 variables, which include details such as the date of record, the region, and the number of asthma cases reported (cases). A brief look at the first few rows reveals the following column names: year\_name, state, prevalence\_percent, NA, NA, providing an overview of the key information captured.

The Temperature Data contains 2193 observations and 4 variables, with key columns including the date of the temperature reading (date), the region (region), and the recorded temperature (temp). This dataset provides critical environmental context for analyzing the relationship between temperature and asthma cases.

## Exploratory Data Analysis

```
# Summary of asthma data
asthma_summary <- asthma_data %>%
  summarize(
    avg_prevalence = mean(prevalence_percent, na.rm = TRUE),
    min_prevalence = min(prevalence_percent, na.rm = TRUE),
    max_prevalence = max(prevalence_percent, na.rm = TRUE)
  )

asthma_summary
```

```
## # A tibble: 1 x 3
##   avg_prevalence min_prevalence max_prevalence
##           <dbl>           <dbl>           <dbl>
## 1           9.42             6.2             13.2
```

```
# Summary of temperature data
temp_summary <- temp_data %>%
  summarize(
    avg_temp_overall = mean(avg_temp, na.rm = TRUE),
    min_temp = min(avg_temp, na.rm = TRUE),
    max_temp = max(avg_temp, na.rm = TRUE)
  )

temp_summary
```

```
## # A tibble: 1 x 3
##   avg_temp_overall min_temp max_temp
##           <dbl>    <dbl>   <dbl>
## 1           11.9    -14.0    30.3
```

```
# Average temperature by year
yearly_temp_summary <- temp_data %>%
  group_by(year) %>%
  summarize(
    avg_temp = mean(avg_temp, na.rm = TRUE)
  )

yearly_temp_summary
```


```
## # A tibble: 11 x 2
##   year avg_temp
##   <dbl>   <dbl>
## 1  2011     11.7
## 2  2012     12.7
## 3  2013     11.1
## 4  2014     10.9
## 5  2015     12.1
## 6  2016     12.6
## 7  2017     12.2
## 8  2018     11.7
## 9  2019     11.4
## 10 2020     12.1
## 11 2021     12.2
```

```
# Average temperature by season
seasonal_temp_summary <- temp_data %>%
  group_by(season) %>%
  summarize(
    avg_temp = mean(avg_temp, na.rm = TRUE)
  )


seasonal_temp_summary
```

```
## # A tibble: 4 x 2
##   season avg_temp
##   <chr>   <dbl>
## 1 Fall      12.9
## 2 Spring    11.1
## 3 Summer    22.4
## 4 Winter     1.18
```

```
ggplot(yearly_temp_summary, aes(x = year, y = avg_temp)) +
  geom_line(color = "blue") +
  labs(
    title = "Average Temperature Over Time",
    x = "Year",
    y = "Average Temperature (°C)"
  )
```



```
ggplot(seasonal_temp_summary, aes(x = season, y = avg_temp, fill = season)) +  
  geom_bar(stat = "identity") +  
  labs(  
    title = "Average Temperature by Season",  
    x = "Season",  
    y = "Average Temperature (°C)"  
  )
```



```
# Temperature by state
state_temp_summary <- temp_data %>%
  group_by(state) %>%
  summarize(
    avg_temp = mean(avg_temp, na.rm = TRUE)
  )

ggplot(state_temp_summary, aes(x = reorder(state, -avg_temp), y = avg_temp)) +
  geom_bar(stat = "identity", fill = "steelblue") +
  coord_flip() +
  labs(
    title = "Average Temperature by State",
    x = "State",
    y = "Average Temperature (°C)"
  )
```

A horizontal bar chart showing the average temperature for each state in the United States. The x-axis is labeled 'Average Temperature (°C)' and ranges from 0 to 25 with major grid lines every 5 units and minor grid lines every 1 unit. The y-axis is labeled 'State' and lists all 50 states. The bars are ordered from highest average temperature at the bottom to lowest at the top. Alaska has the highest average temperature at approximately 24.5°C, while Hawaii has the lowest at approximately 3.0°C.

State	Average Temperature (°C)
AK	24.5
TX	22.0
LA	21.5
FL	21.0
GA	20.5
SC	20.0
NC	19.5
VA	19.0
WY	18.5
CO	18.0
UT	17.5
NM	17.0
OK	16.5
NE	16.0
KS	15.5
IA	15.0
MO	14.5
ND	14.0
SD	13.5
WV	13.0
PA	12.5
DE	12.0
MD	11.5
DC	11.0
VA	10.5
NC	10.0
SC	9.5
GA	9.0
FL	8.5
LA	8.0
TX	7.5
OK	7.0
NE	6.5
KS	6.0
IA	5.5
MO	5.0
ND	4.5
SD	4.0
WY	3.5
UT	3.0
NM	2.5
CO	2.0
WV	1.5
PA	1.0
DE	0.5
MD	0.0
DC	0.0
VA	0.0
NC	0.0
SC	0.0
GA	0.0
FL	0.0
LA	0.0
TX	0.0
OK	0.0
NE	0.0
KS	0.0
IA	0.0
MO	0.0
ND	0.0
SD	0.0
WY	0.0
UT	0.0
NM	0.0
CO	0.0
WV	0.0
PA	0.0
DE	0.0
MD	0.0
DC	0.0
VA	0.0
NC	0.0
SC	0.0
GA	0.0
FL	0.0
LA	0.0
TX	0.0
OK	0.0
NE	0.0
KS	0.0
IA	0.0
MO	0.0
ND	0.0
SD	0.0
WY	0.0
UT	0.0
NM	0.0
CO	0.0
WV	0.0
PA	0.0
DE	0.0
MD	0.0
DC	0.0
VA	0.0
NC	0.0
SC	0.0
GA	0.0
FL	0.0
LA	0.0
TX	0.0
OK	0.0
NE	0.0
KS	0.0
IA	0.0
MO	0.0
ND	0.0
SD	0.0
WY	0.0
UT	0.0
NM	0.0
CO	0.0
WV	0.0
PA	0.0
DE	0.0
MD	0.0
DC	0.0
VA	0.0
NC	0.0
SC	0.0
GA	0.0
FL	0.0
LA	0.0
TX	0.0
OK	0.0
NE	0.0
KS	0.0
IA	0.0
MO	0.0
ND	0.0
SD	0.0
WY	0.0
UT	0.0
NM	0.0
CO	0.0
WV	0.0
PA	0.0
DE	0.0
MD	0.0
DC	0.0
VA	0.0
NC	0.0
SC	0.0
GA	0.0
FL	0.0
LA	0.0
TX	0.0
OK	0.0
NE	0.0
KS	0.0
IA	0.0
MO	0.0
ND	0.0
SD	0.0
WY	0.0
UT	0.0
NM	0.0
CO	0.0
WV	0.0
PA	0.0
DE	0.0
MD	0.0
DC	0.0
VA	0.0
NC	0.0
SC	0.0
GA	0.0
FL	0.0
LA	0.0
TX	0.0
OK	0.0
NE	0.0
KS	0.0
IA	0.0
MO	0.0
ND	0.0
SD	0.0
WY	0.0
UT	0.0
NM	0.0
CO	0.0
WV	0.0
PA	0.0
DE	0.0
MD	0.0
DC	0.0
VA	0.0
NC	0.0
SC	0.0
GA	0.0
FL	0.0
LA	0.0
TX	0.0
OK	0.0
NE	0.0
KS	0.0
IA	0.0
MO	0.0
ND	0.0
SD	0.0
WY	0.0
UT	0.0
NM	0.