

A Datasets and evaluation

Training cohort

44 patients · 352 FCS
7 targets · 8 response channels

Nested patient holdout
Training-only action selection
Once-only outer evaluation





Stanford transport cohort

9 patients · 27 FCS
2 scored targets

Training-frozen policies
No transport refit or recalibration

B Training cohort



95% descriptive percentile intervals
50,000 whole-patient resamples


C Exact channel selections

Selected outer folds / 44

BCR-Crosslink	0	41	0	0	1	0	2	0
BEZ-235	0	0	40	3	0	0	1	0
Dasatinib	0	0	1	42	0	1	0	0
IL-7	0	0	38	5	0	0	1	0
Pervanadate	0	43	0	0	0	0	1	0
TSLP	0	0	0	43	0	1	0	0
Tofacitinib	0	0	2	41	0	0	1	0
pS6								
p4EBP1								
pSTAT5								
pSYK								
pPLCy2								
PAKT								
pERK1/2								
pIKAROS								

Shared: pSYK 43/44; pERK1/2 1/44


D Patient-deletion sensitivity



Points: one patient removed
Dashed: full-cohort contrast

● C0 - Shared
■ C0 - Exact
▲ Shared - Exact

E Stanford transport cohort



Training-frozen scaling; points only
Dasatinib: Shared = Exact (pSYK)
IL-7: Shared pSYK; Exact pSTAT5

● C0 ■ Shared ▲ Exact

F Observed ordering shift

Training cohort

Shared < C0 < Exact

Stanford transport cohort

Exact < Shared < C0

Cohort-specific realized
point estimates

No population-optimum or
site-interaction claim